

(19) World Intellectual Property
Organization
International Bureau



Rec'd PCT TO 19 SEP 2005

10/5498049 804



(43) International Publication Date
30 September 2004 (30.09.2004)

PCT

(10) International Publication Number
WO 2004/083457 A1

(51) International Patent Classification⁷: C12Q 1/68

(21) International Application Number:
PCT/CA2004/000424

(22) International Filing Date: 19 March 2004 (19.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/455,550 19 March 2003 (19.03.2003) US

(71) Applicant (for all designated States except US): THE
UNIVERSITY OF BRITISH COLUMBIA [CA/CA];
University-Industry Liaison Office, 103-6190 Agronomy
Road, Vancouver, British Columbia V6T 1Z3 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RUSSELL, James,
A. [CA/CA]; 1963 West 35th Avenue, Vancouver, British
Columbia V6M 1H8 (CA). WALLEY, Keith, R. [CA/CA];
4172 Coventry Way, North Vancouver, British Columbia
V7N 4M9 (CA).

(74) Agents: ROBINSON, J., Christopher et al.; Smart &
Biggar, Box 11560, Vancouver Centre, Suite 2200, 650
West Georgia Street, Vancouver, British Columbia V6B
4N8 (CA).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PLASMINOGEN ACTIVATOR INHIBITOR-1 (PAI-1) HAPLOTYPES USEFUL AS INDICATORS OF PATIENT OUTCOME

(57) Abstract: The invention provides methods and kits for obtaining a prognosis for a patient having or at risk of developing an inflammatory condition. The method generally comprises determining a Plasminogen Activator Inhibitor-1 (PAI-1) genotype of a patient for one or more polymorphisms in the PAI-1 gene of the patient, comparing the determined genotype with known genotypes for the polymorphism that correspond with the ability of the patient to recover from the inflammatory condition and identifying patients based on their prognosis. PAI-1 genotype screening may be useful in identifying patients who would benefit from increased monitoring by healthcare professionals, and/or possible therapeutic intervention, if the patient were to develop inflammation due to systematic inflammation response syndrome (SIRS), bacterial infection, bacteraemia, sepsis, septic shock, organ dysfunction, and trauma. The invention also provides for methods of identifying other polymorphisms that correspond with the ability of the patients to recover from the inflammatory condition.

BEST AVAILABLE COPY

WO 2004/083457 A1

PLASMINOGEN ACTIVATOR INHIBITOR-1 (PAI-1) HAPLOTYPES USEFUL AS INDICATORS OF PATIENT OUTCOME

RELATED APPLICATION DATA

- 5 This application relates to U.S. provisional application No. 60/455,550 filed March 19, 2003, which is incorporated herein by reference.

FIELD OF THE INVENTION

- 10 The present invention relates to the field of assessment and treatment of patients and more specifically, to the determination of a patients genotype for one or more polymorphisms to determine the patient's prognosis and to determine therapy.

BACKGROUND OF THE INVENTION

- Systemic inflammatory response syndrome (SIRS) is characterized by increased inflammation (anti-inflammatory processes), increased coagulation (anti-coagulant processes), and decreased fibrinolysis (SUFFREDINI AF et al. *New England Journal of Medicine* (1989) 320(18):1165-72; HESSELVIK JF et al. *Critical Care Medicine* (1989) 17(8):724-33; PHILIPPE J et al. *Thrombosis & Haemostasis* (1991) 65(3):291-5; LORENTE JA et al. *Chest* (1993) 103(5):1536-42; FOURRIER F et al. *Chest* (1992) 101(3):816-23; MESTERS RM et al. (1996) 88(3):881-6.).
- 20 Septic and non-septic stimuli such as bacterial endotoxin and cardiopulmonary bypass (CPB), are implicated in the activation of the coagulation system and trigger a systemic inflammatory response syndrome (SIRS). Plasminogen-activator-inhibitor-1 (PAI-1) is associated with decreased fibrinolysis (BINDER BR et al. *News in Physiological Sciences* (2002) 17:56-61). PAI-1 levels are reported increased in patients with sepsis (MESTERS RM et al. *Thrombosis & Haemostasis*. (1996) 75(6):902-7; PRALONG G et al. *Thrombosis & Haemostasis*. (1989) 61(3):459-62), in meningococemia (HERMANS PW et al. *Lancet*. (1999) 354(9178):556-60), and in trauma (MENGENS T et al. *Lancet* (2001) 357(9262):1096-7). Furthermore, increased PAI-1 levels are associated with decreased survival and increased organ dysfunction (MESTERS RM et al. (1996); PRALONG G et al. (1989)).

30

The human PAI-1 gene maps to chromosome 7 q21-q22 and extends over about 12kb. A representative *Homo sapiens* PAI-1 gene sequence is listed in GenBank under accession number AF386492 (shown herein as SEQ ID NO.: 1). DAWSON et al. (*Journal of Biological Chemistry*

(1993) 268(15):10739-45) identified an insertion/deletion polymorphism (4G/5G) at position – 675 of the PAI-1 promoter sequence, which corresponds to position 837 of SEQ ID NO.: 1. The 4G allele is a single base pair deletion promoter polymorphism of the PAI-1 gene and is associated with increased protein levels of PAI-1 (DAWSON SJ et al. (1993); DAWSON SJ et al. *Arteriosclerosis & Thrombosis* (1991) 11(1):183-90). The 4G allele of this single nucleotide polymorphism (SNP) is associated with increased risk of deep venous thrombosis (SEGUI R et al. *British Journal of Haematology* (2000) 111(1):122-8), stroke (HINDORFF LA et al. *Journal of Cardiovascular Risk* (2002) 9(2):131-7), acute myocardial infarction (BOEKHOLDT SM et al. *Circulation* (2001) 104(25):3063-8; ERIKSSON P et al. *PNAS* (1995) 92(6):1851-5.), late lumen loss after coronary artery stent placement (ORTLEPP JR et al. *Clinical Cardiology* (2001) 24(9):585-91), and sudden cardiac death (ANVARI A et al. *Thrombosis Research* (2001) 103(2):103-7; MIKKELSSON J et al. *Thrombosis & Haemostasis* (2000) 84(1):78-82). In the critically ill, the 4G allele is also associated with decreased survival in patients who have had severe trauma (MENGES T et al. *Lancet* (2001) 357(9262):1096-7) and patients who had meningococemia (HERMANS PW et al. *Lancet*. (1999) 354(9178):556-60) as well as increased risk of shock in patients who had meningococemia (WESTENDORP RG et al. *Lancet* (1999) 354(9178):561-3). The PAI-1 4G genotype has also been associated with adverse patient outcomes ((MENGES et al. (2001); HERMANS et al. (1999); WESTENDORP RG et al. (1999); ENDLER G et al. *British Journal of Haematology* (2000) 110(2):469-71; GARDEMANN A et al. *Thrombosis & Haemostasis* (1999) 82(3):1121-6; HOOPER WC et al. *Thrombosis Research* (2000) 99(3):223-30; JONES K et al. *European Journal of Vascular & Endovascular Surgery* (2002) 23(5):421-5; HARALAMBOUS E. et al. *Crit Care Med* (2003) 31(12):2788-93; and ROEST M et al. *Circulation* (2000) 101(1):67-70). The 4G/4G genotype of PAI-1 was associated with PAI-1 levels in patients suffering from acute lung injury (RUSSELL JA *Crit Care Med*. (2003) 31(4):S243-S247).

SUMMARY OF THE INVENTION

This invention is based in part on the surprising discovery that the Plasminogen-Activator-Inhibitor-1 (PAI-1) 4G/5G promoter polymorphism is not as accurate a predictor of patient outcome as the SNP at position 12580 of SEQ ID NO:1. Furthermore, various other PAI-1 polymorphisms in total linkage disequilibrium (LD) with SNP 12580 are identified as being useful for patient screening, as an indication of patient outcome, or for prognosis for recovery from an inflammatory condition or to tailor therapy for patients with inflammatory conditions or

for the selection of therpaies for patients based on their PAI-1 12580 genotype and hapoltype. Surprisingly, the SNPs identified herein as predictors of patient outcome also correspond with lower PAI-1 levels.

- 5 In accordance with one aspect of the invention, methods are provided for obtaining a prognosis for a patient having or at risk of developing an inflammatory condition, the method comprising determining a genotype including one or more polymorphism sites in the PAI-1 gene for the patient, wherein said genotype is indicative of an ability of the patient to recover from an inflammatory condition and provided the genotype selected does not correspond to only the 837
10 position of SEQ ID NO.: 1.

In accordance with another aspect of the invention, methods are provided for obtaining a prognosis for a patient having or at risk of developing an inflammatory condition, the method comprising determining a genotype of one or more polymorphism sites in the PAI-1 gene for the
15 patient selected from the polymorphism sites corresponding to positions 664, 2037, 2362, 2852, 5834, 5878, 7343, 13605, 7365, 7729, 7771, 12750, 4588, 5404, 5686, 5984, 11312, 2846, 6821, 9759, 10381, 7365, 7729, 7771, 12750, 4588, 5404, 5686, 5984, 11312, 5645, 7121, 7437, 8070, 8406, 9463, 9466, 12219, 12580, 13889, 14440, wherein said genotype is indicative of an ability of the patient to recover from an inflammatory condition.

20 In accordance with one aspect of the invention, methods are provided for obtaining a prognosis for a patient having or at risk of developing an inflammatory condition, the method comprising determining a genotype including one or more polymorphism sites in the PAI-1 gene for the patient, wherein said genotype is indicative of an ability of the patient to recover from an
25 inflammatory condition and the polymorphism is selected from the region defined by positions 1581 to 14544 of SEQ ID NO.: 1.

The polymorphism site may correspond to position 12580 of SEQ ID NO.: 1 or a polymorphism site linked thereto. Alternatively, the polymorphism site may correspond to positions 5645, 7121,
30 7437, 8070, 8406, 9463, 9466, 12219, 13889 or 14440 in SEQ ID NO: 1.

Genotype may also be determined at a combination of multiple polymorphism sites, the combination being selected from the group of positions corresponds to SEQ ID NO:1 consisting

of:

664 and 2037;

664 and 2362;

664 and 2852;

5 664 and 5834;

837 and 2037;

837 and 2362;

837 and 2852;

837 and 5834;

10	one of 5878 and one of	7365	and one of	4588
	7343	7729		5404
	13605	7771		5686
		12750		5984
				11312; and
15	one of 2846 and 10381	and one of 7365	and one of	4588
	6821	7729		5404
	9759	7771		5686
		12750		5984
				11312.

20

Alternatively, genotypes may be determined for polymorphism located within the 1581-14544 of SEQ ID NO:1, to exclude polymorphisms within the promoter region. Preferably, genotypes would be determined for polymorphisms selected from the group consisting of 5645, 7121, 7437, 8070, 8406, 9463, 9466, 12219, 12580, 13889, or 14440 or SEQ ID NO:1.

25

In accordance with another aspect of the invention, methods are provided for further comparing the genotype so determined with known genotypes, which are indicative of a prognosis for recovery from the same inflammatory condition as for the patient or another inflammatory condition.

30

The genotype of the patient may be indicative of a decreased likelihood of recovery from an inflammatory condition or indicative of a prognosis of severe cardiovascular or respiratory dysfunction in critically ill patients. Furthermore, such a genotype may be selected from the

group of single polymorphism sites and combined polymorphism sites consisting of:

5 5645 T;
7121 G;
7437 T;
8070 A;
8406 C;
9463 G;
9466 T;
10 12219 C;
12580 G;
13889 C;
14440 A;
15 664 A and 2037 T;
664 A and 2362 -;
664 A and 2852 A;
664 A and 5834 A;
837 - and 2037 T;
837 - and 2362 -;
20 837 - and 2852 A; and
837 - and 5834 A.

The genotype of the patient may be indicative of an increased likelihood of recovery from an inflammatory condition or indicative of a prognosis of less severe cardiovascular or respiratory
25 dysfunction in critically ill patients. Furthermore, such a genotype may be selected from the group of single polymorphism sites and combined polymorphism sites consisting of:

30 5645 C;
7121 A;
7437 C;
8070 G;
8406 T;
9463 A;
9466 C;
12219 T;
35 12580 T;
13889 T;
14440 G;
one of 5878 G and one of 7365 T and one of 4588 T
7343 G 7729 + 5404 G
40 13605 A 7771 A 5686 A
12750 A 5984 A
11312 A; and
one of 2846 A and 10381 T and one of 7365 T and one of 4588 T
45 6821 T 7729 + 5404 G
9759 G 7771 A 5686 A

12750 A

5984 A
11312 A.

5 In accordance with another aspect of the invention, methods are provided for identifying a polymorphism in a PAI-1 gene sequence that correlates with patient prognosis. Where the method comprises obtaining PAI-1 gene sequence information from a group of patients, identifying a site of at least one polymorphism in the PAI-1 gene, determining genotypes at the site for individual patients in the group, determining an ability of individual patients in the group to recover from the inflammatory condition and correlating genotypes determined with patient
10 with potential therapies.

The correlation procedure may be repeated on a patient population of sufficient size to achieve a statistically significant correlation.

15 The methods may further comprise steps of obtaining PAI-1 gene sequence of the patient or obtaining a nucleic acid sample from the patient. The determining of genotype may be performed on a nucleic acid sample from the patient.

20 Where the genotype of the patient corresponding to the nucleotide in position 12580, is G, the prognosis may be indicative of a decreased likelihood of recovery from an inflammatory condition or of severe cardiovascular or respiratory dysfunction in critically ill patients.

Where the genotype of the patient corresponding to the nucleotide in position 12580, is T, the prognosis may be indicative of a increased likelihood of recovery from an inflammatory condition
25 or of less severe cardiovascular or respiratory dysfunction in critically ill patients.

The inflammatory condition may be selected from the group consisting of: sepsis, septicemia, pneumonia, septic shock, systemic inflammatory response syndrome (SIRS), Acute Respiratory Distress Syndrome (ARDS), acute lung injury, infection, pancreatitis, bacteremia, peritonitis,
30 abdominal abscess, inflammation due to trauma, inflammation due to surgery, chronic inflammatory disease, ischemia, ischemia-reperfusion injury of an organ or tissue, tissue damage due to disease, tissue damage due to chemotherapy or radiotherapy, and reactions to ingested, inhaled, infused, injected, or delivered substances, glomerulonephritis, bowel infection,

opportunistic infections, and for patients undergoing major surgery or dialysis, patients who are immunocompromised, patients on immunosuppressive agents, patients with HIV/AIDS, patients with suspected endocarditis, patients with fever, patients with fever of unknown origin, patients with cystic fibrosis, patients with diabetes mellitus, patients with chronic renal failure, patients with bronchiectasis, patients with chronic obstructive lung disease, chronic bronchitis, emphysema, or asthma, patients with febrile neutropenia, patients with meningitis, patients with septic arthritis, patients with urinary tract infection, patients with necrotizing fasciitis, patients with other suspected Group A streptococcus infection, patients who have had a splenectomy, patients with recurrent or suspected enterococcus infection, other medical and surgical conditions associated with increased risk of infection, Gram positive sepsis, Gram negative sepsis, culture negative sepsis, fungal sepsis, meningococcemia, post-pump syndrome, cardiac stun syndrome, myocardial infarction, stroke, congestive heart failure, hepatitis, epiglottitis, E. coli 0157:H7, malaria, gas gangrene, toxic shock syndrome, mycobacterial tuberculosis, Pneumocystic carinii, pneumonia, Leishmaniasis, hemolytic uremic syndrome/thrombotic thrombocytopenic purpura, Dengue hemorrhagic fever, pelvic inflammatory disease, Legionella, Lyme disease, Influenza A, Epstein-Barr virus, encephalitis, inflammatory diseases and autoimmunity including Rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, inflammatory bowel disease, idiopathic pulmonary fibrosis, sarcoidosis, hypersensitivity pneumonitis, systemic vasculitis, Wegener's granulomatosis, transplants including heart, liver, lung kidney bone marrow, graft-versus-host disease, transplant rejection, sickle cell anemia, nephrotic syndrome, toxicity of agents such as OKT3, cytokine therapy, and cirrhosis.

The determining of a genotype may comprise one or more of: restriction fragment length analysis; sequencing; hybridization; oligonucleotide ligation assay; ligation rolling circle amplification; 5' nuclease assay; polymerase proofreading methods; allele specific PCR; and reading sequence data.

In accordance with another aspect of the invention, there is provided a kit for determining a genotype at a defined nucleotide position within a polymorphism site in a PAI-1 gene sequence from a patient to provide a prognosis of the patient's ability to recover from an inflammatory condition, the kit comprising, in a package a restriction enzyme capable of distinguishing alternate nucleotides at the polymorphism site or a labeled oligonucleotide having sufficient complementary to the polymorphism site and capable of distinguishing said alternate nucleotides.

The alternate nucleotides may correspond to position 12580 of SEQ ID NO: 1, position 10 of SEQ ID NO: 2 or to a polymorphism linked thereto. The alternate nucleotides may also correspond to one or more of positions 5645, 7121, 7437, 8070, 8406, 9463, 9466, 12219, 13889 or 14440 of SEQ ID NO: 1.

The kit comprising a restriction enzyme may also comprise an oligonucleotide or a set of oligonucleotides suitable to amplify a region surrounding the polymorphism site, a polymerization agent and instructions for using the kit to determine genotype.

In accordance with another aspect of the invention, methods are provided for determining patient prognosis in a patient having or at risk of developing an inflammatory condition, the method comprising detecting the identity of one or more polymorphisms in the PAI-1 promoter region, wherein the identity of said one or more polymorphisms is indicative of the ability of the patient to recover from the inflammatory condition.

In accordance with another aspect of the invention, methods are provided for patient screening, comprising the steps of (a) obtaining PAI-1 gene sequence information from a patient, and (b) determining the identity of one or more polymorphisms in the promoter region, wherein the one or more polymorphisms may be indicative of the ability of a patient to recover from an inflammatory condition.

In accordance with another aspect of the invention methods are provided for patient screening whereby the method includes the steps of (a) selecting a patient based on risk of developing an inflammatory condition or having an inflammatory condition, (b) obtaining PAI-1 gene sequence information from the patient and (c) detecting the identity of one or more polymorphisms in the PAI-1 gene, wherein the polymorphism is indicative of the ability of a patient to recover from an inflammatory condition.

In accordance with another aspect of the invention methods are provided for selecting a group of patients to determine the efficacy of a candidate drug known or suspected of being useful for the treatment of an inflammatory condition, the method including determining a genotype for one or more polymorphism sites in the plasminogen-activator-inhibitor-1 gene for each patient, wherein

said genotype is indicative of the patient's ability to recover from the inflammatory condition and sorting patients based on their genotype, provided that the polymorphism site is not solely a polymorphism at a site corresponding to position 837 of SEQ ID NO.: 1.

- 5 The method may include the additional step of comparing patient response to the candidate drug based on genotype of the patient. Response to the candidate drug may be decided by determining each patient's ability to recover from the inflammatory condition.

The above sequence positions refer to the sense strand of the PAI-1 gene. It will be obvious to a person skilled in the art that analysis could be conducted on the anti-sense strand to determine patient outcome.

More severe patient outcome or prognosis may be correlated with lower PAI-1 expression or conversely an indication of less severe patient outcome or prognosis may be correlated with higher PAI-1 expression, which is the opposite of what would be expected.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. shows a haplotype map of plasminogen activator inhibitor-1 (PAI-1) in Caucasians, where the columns indicate polymorphic loci and rows indicate distinct haplotypes. Light cells represent rare alleles, and dark cells represent common alleles. Polymorphisms that provide the same information as the 12580G allele were chosen based on this haplotype map. These polymorphisms provide the same information as the 12580G allele.

25 Figure 2. shows the PAI-1 haplotype clades in an unrooted phylogenetic tree indicating the positions of haplotypes from Figure 1. Figure 2 was drawn using MEGA II where line length reflects relative evolutionary distance and each clade is circled.

Figure 3. shows hospital survival for each of the haplotype clades of the PAI-1 gene. Error bars indicate 95% confidence intervals. Haplotypes containing the 4G allele are made up of 2 separate haplotype clades defined by: the 13985 T allele and the 12580 G allele. The group mean, excluding the 4G haplotype clade containing the 12580 G allele, is illustrated by the horizontal line. The mean and 95% confidence interval for the haplotype clade defined by the 12580 G allele falls below the mean of the rest of the

group. In contrast, the 4G haplotype clade containing the 13985 T allele has the highest hospital survival of any haplotype. Data from patients were included twice, once for each haplotype that they possessed (homozygotes for a haplotype were included twice in that haplotype), so that the number of haplotypes (numbers within the data bars) add up to 2 times the total number of patients (235).

Figure 4. shows Days Alive and Free (DAF) of organ system failure over the 28 day study period by genotype for the PAI-1 12580 allele.

Figure 5 shows days alive and free of fulfillment of 2, 3 or 4 out of 4 (2/4, 3/4 or 4/4) systemic inflammatory response syndrome (SIRS) criteria by PAI-1 12580 genotype. The SIRS criteria are: 1) tachycardia (heart rate >100 bpm in the absence of beta blockers), 2) tachypnea (respiration rate > 20 breaths/min) or mechanically ventilated, 3) fever (temperature > 38°C) or hypothermia (temperature < 35.5°C), and 4) leukocytosis (total leukocyte count > 11,000/ μ L), using the guidelines established in the American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference:

definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis (*Critical Care Medicine*, (1992) 20(6):864-74). Each day that a patient was in the ICU the presence or absence of each SIRS criteria was scored, such that the total number of SIRS criteria fulfilled on each day in the ICU was tallied. Since fulfillment of SIRS criteria indicates poor outcome, more days alive and free of fulfillment of SIRS criteria indicates a better outcome.

Figure 6. shows the occurrence of Sepsis, and Septic Shock by genotype for the PAI-1 12580 allele.

Figure 7A. shows days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 12580 genotype (GG, GT and TT separately) in 587 critically ill Caucasians with the systemic inflammatory response syndrome (SIRS).

Figure 7B. shows days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 12580 genotype (GG/GT vs TT) in 587 critically ill Caucasians with SIRS.

Figure 8A. shows days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 4G/5G (837) genotype (4G/4G, 4G/5G and 5G/5G separately) in 314 critically ill Caucasians with the systemic inflammatory response syndrome (SIRS).

Figure 8B. shows days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 4G/5G (837) allele (any 4G vs 5G/5G) in 314 critically ill Caucasians with SIRS.

Figure 9A. shows days alive and free of acute lung injury (ALI) and P_aO_2/F_iO_2 (P/F) ratio greater than 300 mmHg by PAI-1 12580 genotype (GG, GT and TT separately) in 587 critically ill Caucasians with SIRS.

Figure 9B. shows days alive and free of acute lung injury and $P_aO_2/F_iO_2 < 300$ mmHg by PAI-1 12580 genotype (GG/GT vs TT) for 587 Caucasians with SIRS.

Figure 10A. shows days alive and free of acute lung injury (ALI) and P_aO_2/F_iO_2 (P/F) ratio greater than 300 mmHg by PAI-1 4G/5G/ (837) genotype (4G/4G, 4G/5G and 5G/5G separately) in 314 critically ill Caucasians with SIRS.

Figure 10B. shows days alive and free of acute lung injury and $P_aO_2/F_iO_2 < 300$ mmHg by PAI-1 4G/5G allele (any 4G vs 5G/5G) in 314 critically ill Caucasians with SIRS.

Figure 11A. shows days alive and free of steroid use by PAI-1 12580 genotype in 587 critically ill Caucasians with SIRS. Patients with 2 copies of the 12580G allele (N = 98) had the fewest days alive and free of steroid use (15 days).

Figure 11B. shows days alive and free of steroid use by PAI-1 12580 genotype (GG/GT vs TT) for 587 critically ill Caucasians with SIRS.

Figure 12B. shows days alive and free of steroid use by PAI-1 4G/5G (837) genotype in critically ill Caucasians with SIRS.

Figure 12B. shows days alive and free of steroid use by PAI-1 4G/5G (837) allele in 314 critically ill Caucasians with SIRS.

Figure 13A. shows mean pre-operative PAI-1 serum levels in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG, GT and TT separately).

Figure 13B. shows mean pre-operative serum PAI-1 levels in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG/GT vs TT).

Figure 14A. shows serum PAI-1 levels measured 4 hours post-operatively in 55 Caucasians patients who underwent cardiopulmonary bypass surgery grouped by PAI-1 4G/5G.

Figure 14B. shows serum PAI-1 levels measured 4 hours post-operatively in 55 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 4G/5G.

Figure 15A. shows mean serum PAI-1 levels of 67 Caucasians who underwent on-pump cardiopulmonary bypass surgery collected at 3 hours post surgery grouped by PAI-1 12580 genotype (TT, GT and GG separately).

Figure 15B. shows mean serum PAI-1 levels of 67 Caucasians who underwent on-pump cardiopulmonary bypass surgery collected 3 hours post-operatively grouped by 12580 GG/GT vs. TT.

Figure 16. shows mean differences in serum PAI-1 levels from pre-operative to 3 hours post-operative in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 GG, GT, and TT.

Figure 17A. shows systemic vascular resistance index (SVRI) 4 hours post-operatively in 66 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG, GT and TT separately).

Figure 17B. shows Systemic vascular resistance index (SVRI) 4 hours post-operatively in 66 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG/GT vs TT).

Figure 18A. shows the systemic vascular resistance index (SVRI) 4 hours post-operatively in 364 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 4G/5G (837) genotype (4G/4G, 4G/5G and 5G/5G separately).

Figure 18B. shows the systemic vascular resistance index (SVRI) 4 hours post-operatively in 364 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 4G/5G (837) allele (any 4G vs 5G/5G).

DETAILED DESCRIPTION OF THE INVENTION

1. Definitions

In the description that follows, a number of terms are used extensively, the following definitions are provided to facilitate understanding of the invention.

"Genetic material" includes any nucleic acid and can be a deoxyribonucleotide or ribonucleotide polymer in either single or double-stranded form.

A "purine" is a heterocyclic organic compound containing fused pyrimidine and imidazole rings, and acts as the parent compound for purine bases, adenine (A) and guanine (G). "Nucleotides" are generally a purine (R) or pyrimidine (Y) base covalently linked to a pentose, usually ribose or

deoxyribose, where the sugar carries one or more phosphate groups. Nucleic acids are generally a polymer of nucleotides joined by 3' 5' phosphodiester linkages. As used herein "purine" is used to refer to the purine bases, A and G, and more broadly to include the nucleotide monomers, deoxyadenosine-5' -phosphate and deoxyguanosine-5' -phosphate, as components of a polynucleotide chain.

A "pyrimidine" is a single-ringed, organic base that forms nucleotide bases, cytosine (C), thymine (T) and uracil (U). As used herein "pyrimidine" is used to refer to the pyrimidine bases, C, T and U, and more broadly to include the pyrimidine nucleotide monomers that along with purine nucleotides are the components of a polynucleotide chain.

A "polymorphic site" or "polymorphism site" or "polymorphism" or "single nucleotide polymorphism site" (SNP site) as used herein is the locus or position within a given sequence at which divergence occurs. A "Polymorphism" is the occurrence of two or more forms of a gene or position within a gene (allele), in a population, in such frequencies that the presence of the rarest of the forms cannot be explained by mutation alone. The implication is that polymorphic alleles confer some selective advantage on the host. Preferred polymorphic sites have at least two alleles, each occurring at frequency of greater than 1%, and more preferably greater than 10% or 20% of a selected population. Polymorphism sites may be at known positions within a nucleic acid sequence or may be determined to exist using the methods described herein. Polymorphisms may occur in both the coding regions and the noncoding regions (for example, upstream promoters) of genes.

A "clade" is a group of haplotypes that are closely related phylogenetically. For example, if haplotypes are displayed on an phylogentic (evolutionary) tree a clade includes all haplotypes contained within the same branch.

As used herein "haplotype" is a set of alleles of closely linked loci on a chromosome that tend to be inherited together; commonly used in reference to the linked genes of the major histocompatibility complex.

As used herein "linkage disequilibrium" is the occurrence in a population of certain combinations of linked alleles in greater proportion than expected from the allele frequencies at the loci.

Numerous other sites have been identified as polymorphisms in the PAI-1 gene, where those polymorphisms are linked to the polymorphism at position 12580 of SEQ. ID NO: 1 and may therefore be indicative of patient prognosis. The following single polymorphism sites and combined polymorphism sites are linked to 12580 of SEQ. ID NO.: 1:

5 5645;
7121;
7437;
8070;
10 8406;
9463;
9466;
12219;
13889;
15 14440;
664 and 2037;
664 and 2362;
664 and 2852;
664 and 5834;
20 837 and 2037;
837 and 2362;
837 and 2852; and
837 and 5834.

25 It will be appreciated by a person of skill in the art that further linked single polymorphism sites and combined polymorphism sites could be determined. The haplotype of PAI-1 can be created by assessing the SNP's of PAI-1 in normal subjects using a program that has an expectation maximization algorithm (i.e. PHASE). A constructed haplotype of PAI-1 may be used to find combinations of SNP's that are in total linkage disequilibrium (LD) with 12580. Therefore, the
30 haplotype of an individual could be determined by genotyping other SNP's that are in total LD with 12580. Linked single polymorphism sites or combined polymorphism sites may also be genotyped for assessing patient prognosis.

The following genotypes for single polymorphism sites and combined polymorphism sites may be
35 indicative of a decreased likelihood of recovery from an inflammatory condition or indicative of severe cardiovascular or respiratory dysfunction in critically ill patients:

5645 T;
7121 G;
7437 T;
40 8070 A;

8406 C;
 9463 G;
 9466 T;
 12219 C;
 12580 G;
 13889 C;
 14440 A;
 664 A and 2037 T;
 664 A and 2362 -;
 664 A and 2852 A;
 664 A and 5834 A;
 837 - and 2037 T;
 837 - and 2362 -;
 837 - and 2852 A; and
 837 - and 5834 A.

Whereas the following genotypes for single polymorphism sites and combined polymorphism sites may be indicative of a increased likelihood of recovery from an inflammatory condition or indicative of less severe cardiovascular or respiratory dysfunction in critically ill patients:

20	5645 C;			
	7121 A;			
	7437 C;			
	8070 G;			
	8406 T;			
25	9463 A;			
	9466 C;			
	12219 T;			
	12580 T;			
	13889 T;			
30	14440 G;			
	one of 5878 G and one of	7365 T	and one of	4588 T
	7343 G	7729 +		5404 G
	13605 A	7771 A		5686 A
		12750 A		5984 A
35	one of 2846 A and 10381 T and one of	7365 T	and one of	11312 A; and
	6821 T and 10381 T	7729 +		4588 T
	9759 G and 10381 T	7771 A		5404 G
		12750 A		5686 A
40				5984 A
				11312 A.

(wherein a "+" represents an insertion and a "-" represents a deletion)

It will be appreciated by a person of skill in the art, that the numerical designations of the positions of polymorphisms within a sequence are relative to the specific sequence. Also the

same positions may be assigned different numerical designations depending on the way in which the sequence is numbered and the sequence chosen, as illustrated by the alternative numbering of equivalent polymorphism in DAWSON *et al.* (1993) above. Furthermore, sequence variations within the population, such as insertions or deletions, may change the relative position and subsequently the numerical designations of particular nucleotides at and around a polymorphism site.

TABLE 1A below is representative of a *Homo sapiens* Plasminogen Activator Inhibitor-1 (PAI-1) or Serine-cysteine Proteinase Inhibitor clade E member 1 (SERPINE1)

gene sequence and comprises a sequence as listed in GenBank under accession number AF386492. The 4G/5G polymorphism of PAI-1 corresponds to the SNP described as -675 G/- using the numbering system of DAWSON *et al.* (1993), which corresponds to position 837 in TABLE 1A (SEQ ID NO: 1). Polymorphism sites shown below in TABLE 1A are bolded and underlined. The polymorphism sites are represented by their major allele (dark cells in Figure 1). The major and minor alleles for each of the primary polymorphism sites of the PAI-1 gene are as follows:

at position 5445 the most common nucleotide (major allele) is **g** and the minor allele is **a**;
at position 837 the most common nucleotide (major allele) is **g** and the minor allele is **c**;
at position 5445 the most common nucleotide (major allele) is **c** and the minor allele is **t**;
at position 7121 the most common nucleotide (major allele) is **a** and the minor allele is **g**;
at position 7437 the most common nucleotide (major allele) is **c** and the minor allele is **t**;
at position 8070 the most common nucleotide (major allele) is **g** and the minor allele is **a**;
at position 8406 the most common nucleotide (major allele) is **t** and the minor allele is **c**;
at position 9463 the most common nucleotide (major allele) is **a** and the minor allele is **g**;
at position 9466 the most common nucleotide (major allele) is **c** and the minor allele is **t**;
at position 12219 the most common nucleotide (major allele) is **t** and the minor allele is **c**;
at position 12580 the most common nucleotide (major allele) is **t** and the minor allele is **g**;
at position 13889 the most common nucleotide (major allele) is **t** and the minor allele is **c**;
at position 14440 the most common nucleotide (major allele) is **g** and the minor allele is **a**.

TABLE 1A

1	1	cgtactggtc	catagcctgt	taggaaccag	gctgcataac	aggaggtgag	tggcaggtga
61		gtgaaatttc	atctgtagtt	acagccactc	ctcatcactc	gcattaccac	cagagctcca
5	121	ctccctgtca	gatcagcggc	ggcattagat	tctcatagga	gctcgaaccc	tattctaaac
	181	tgttcatgtg	agggatctag	gttgcaagct	ccctatgaga	atctaatagc	tgatgatctg
	241	tcacggctctc	ccatcacccc	tagatgggac	catctagttg	caggaaaaca	agctcagggc
	301	tccactgat	tctacacgat	ggtgaattgt	ggaattatct	cattatatat	attacaatgt
	361	aataataata	gaaataaagc	acacaataaa	tgtaatgtgc	ttgaatcatc	ccgaaaccat
10	421	cccaccctgg	tctgtgaaaa	aattgtcttc	catgaaacca	gtccctgggtg	ccaaaaacgt
	481	tgaggaccac	tgtctccacag	aatctatcgg	tactcttcc	tcccctcacc	cccttgccct
	541	aaaagcacac	cctgcaaac	tgccatgaat	tgacactctg	tttctatccc	ttttcccctt
	601	gtgtctgtgt	ctggaggaag	aggataaagg	acaagctgcc	ccaagtctta	gcgggcagct
	661	cgagggaagtg	aaacttacac	gttggtctcc	tgtttcctta	ccaagctttt	accatggtaa
15	721	ccctgggtcc	cgttcagcca	ccaccacccc	accagcaca	cctccaacct	cagccagaca
	781	aggttggtga	cacaagagag	ccctcagggg	cacagagaga	gtctggacac	gtgggggagt
	841	cagccgtgta	tcacgcggag	cggccgggca	catggcaggg	atgagggaaa	gaccaagagt
	901	cctctgttgg	gcccaggtcc	tagacagaca	aaacctagac	aatcacgtgg	ctggctgcat
	961	gccctgtggc	tgttgggctg	ggcccaggag	gagggagggg	cgtcttttcc	tggaggtggt
20	1021	ccagagcacc	gggtggacag	ccctggggga	aaacttcac	gttttgatgg	aggttatctt
	1081	tgataactcc	acagtgacct	ggttcgcca	aggaaaagca	ggcaacgtga	gctgtttttt
	1141	ttttctccaa	gctgaacact	aggggtccta	ggcttttttg	gtcaccgggc	atggcagaca
	1201	gtcaacctgg	caggacatcc	gggagagaca	gacacaggca	gagggcagaa	aggtcaaggg
	1261	aggttctcag	gccaaggcta	ttgggggttg	ctcaattgtt	cctgaatgct	cttacacacg
25	1321	tacacacaca	gagcagcaca	cacacacaca	cacacacatg	cctcagcaag	tcccagagag
	1381	ggaggtgtcg	agggggaccc	gctggctgtt	cagacggact	cccagagcca	gtgagtgggt
	1441	ggggctggaa	catgagttca	tctatttcct	gcccacatct	ggtataaaaag	gaggcagtgg
	1501	cccacagagg	agcacagctg	tgtttggctg	cagggccaag	agcgtgttca	agaagacca
	1561	cacgcccccc	tccagcagct	gaattcctgc	agctcagcag	ccgcccgcag	agcaggacga
30	1621	acgcgcaatc	gcaaggcacc	tctgagaact	tcaggtagga	gaaaagcaaa	ctccctccaa
	1681	cctcttactt	cgggcttaag	gcagagaact	cgcctcccca	gaatctcttc	cctccatgat
	1741	ccccgcctat	tcctctatct	tcttttcttc	ggacctgcag	ccttgggctg	accctgccct
	1801	aggggtgact	gcaggagagc	agggaggatg	gtcaggcgct	accaacaacc	ccatcaccca
	1861	gtaacaagaa	ccttgactct	ctcagtcctt	ctgcatcaag	acacttaccc	atttcccacc
35	1921	tcatgcctgc	taacttgaat	gaaacaatcg	ctgggaaagc	attaagagaa	ttaaggctgg
	1981	gcactgtggc	tcatgcctgt	aatcccagca	ctttgtgagg	ctgaggcagg	cagataactt
	2041	gagcccagga	gtttgagacc	agcctgggca	acatggcaaa	accctgctct	cccaaaaaaa
	2101	tacaaaaatt	agctgggcgt	gctgggtgtg	ctgtattccc	agctacttgg	gaggctgagg
	2161	tgggaggatt	gcttcagctg	gggaggcgga	ggctgcaggg	agccaagact	gagccattgc

2221 acccagcctg ggtgacagag caagaccctg tctctaaaaa tgaatgaaag gaaggaagaa
2281 agagagagaa agagagagag gaaagaagga aggaagtaaa gaagaaagaa agaaagaaag
2341 aggaaagagg aagaaagaaa gaaaagaaag aaaagaaaag aaagcaaatt taaagcttat
2401 gcaaatacaa gatgttgta taattgataa ttgagtctgg gctaaattcc ccctgggctg
5 2461 caaaggcaga gagtggtaat gacttctcac ctgcttttct tctaaggctt ttttacggga
2521 cacagaggga agggagatgg actggattcc aagattccca cagggcaaga tgggcgaaga
2581 ctccctgcc a ctgccgggg ataagtcagt ctgagtgaga cggagtggga tgggcttaga
2641 acctgaacat gtcattggtct ctccctgcac cttgccctag tgttcaacta ccacctgott
2701 gcaggaaaca agaagagcag ggcccacagc tggccagctc ccctccctc ccgctgtct
10 2761 tccagaacga ttccttcacc agccctcttt ccattgctct aggatgcaga tgtctccagc
2821 cctcacctgc ctagtcttg gcctggccct tgtctttggt gaagggtctg ctgtgcacca
2881 tccccatcc tacgtggccc acctggcctc agacttcggg gtgaggggtgt ttcagcaggt
2941 ggcgaggcc tccaaggacc gcaacgtggt tttctcacc tatgggggtg cctcggtgtt
3001 ggccatgctc cagctgacaa caggaggaga aaccagcag cagattcaag cagctatggg
15 3061 attcaagatt gatggtgagc caggggacac caggggaggt ggggtggcatg cagaacagac
3121 ctaccagaag ccaaggaaag gctggctctg gcttagccga gccaaagccc atacagctgt
3181 gctgcagggg ccaccccatc ttcttccac tactctcaa gtcactggac ccttgaatct
3241 ccaagggtgt ctgaccagta gatttaccgc ttattcacca ccgtgtgatc ttaacctcgt
3301 taagtttgcc catctacaaa atgaggatta tttgctgtcc taaagaattc atgagccggg
20 3361 cgcggtggct caaacgcctg taatcccagc actttgggag gccaaaggcg gcggatcatg
3421 aggtcaggag atcaagacca tcctggctaa cacagtgaag ctccatctct actaaaaata
3481 caaaaaaat tagccaggcg tgggtggcagg cgctgtagt ccagctact cgggaggctg
3541 aggcaggaga atggcatgaa ccaggaggc agagcttgca gtgagctgag atcgtgccac
3601 tgcactccag cctgggagc agagagagac tccgtctcaa aaaaaaaat taaaagaatt
25 3661 catggaatta cacttgtaaa atacttagca tagccatcac tataggaaaa aaatctaagg
3721 ccaggcacag tgcctcatgc ctgtaattct agcactttcg gagtttgagg caggaggatc
3781 acccaaggct aggagttcaa ggccagcctg ggcaatacgg tgaaacccc tctctaataa
3841 aaatataaaa attagtctga tgaggtggtg cacctgtaat ccagctact caggaagctg
3901 agacacaaga atcactttaa cccgggaggt ggaggtggca gtgagctgag atcacaccat
30 3961 tgcactccag cctgggtgac agagtgcagc ctgtcaaaaa aaagaaaaga aagagagaga
4021 gagagagaag agagagaaag aaagaagaag aaagaaagaa agagagagag agaaagaaag
4081 aaagaaacaa agaaacaaag aaagaaagga aaagaaaaaa aaaactaagg ccaggcaagg
4141 tggcttatga ctgtaatttc agcactttgg aagattgagg caggaggatc acttgaggcc
4201 agaagttcga gacaagactg agcaacaggg agaccctgc ctctacaaaa aaatttacia
35 4261 attagccaga tgtggtgaca catacctgta gtcccaacta ctcaggaggc tgaggtggga
4321 ggatggcttg agcccaggag ctggaggctg cagtgcagta tgattgtacc actgcacttc
4381 agcctgggca acaagggaa gccctgtctg aaaaaaaaaa aaaaagaaaa agaagaagaa
4441 agaaaatatt tagggttcac ccaggaggca gaggttgag taagctgaca tcgcccatt
4501 gcactccagc ctgggagaca agagcaaac tccaactcaa aaaaaaaaaa aaaaaaaaaa
40 4561 caggaagaaa atatttaggg ttcataactt aagaacagag aaaaatatc tagcccaag

4621 aaaggggttg gatctgagac ttttgaagaa aggaaggaga tacagaaaag agatttcac
 4681 ctggaatgaa atctccctcc agagagccct gggaaagcac ggtagcccc atccatcaga
 4741 gtggagcccc ttgtggggga agtgggctcg gctgggaacc ctcaattcag cataagcctc
 4801 acatgtctc tcctctctgt cccggtgcag acaagggcat ggccccgcc ctccggcatc
 5 4861 tgtacaagga gctcatggg ccatggaaca aggatgagat cagcaccaca gacgcgatct
 4921 tcgtccagcg ggatctgaag ctgggtccagg gcttcatgcc ccacttcttc aggctgttcc
 4981 ggagcacggt caagcaagtg gacttttcag aggtggagag agccagattc atcatcaatg
 5041 actgggtgaa gacacacaca aaaggtgagc aggcaggga aggaaacca tttcctgggc
 5101 ctcaagagaa agggaatttg gaaataaatc cacatatccc agttgggtgc agtagttcac
 10 5161 acctgtaatc ccagcccaac actttgggag gtctaggcga gaggaaggct tgaggcctgg
 5221 agtttgagac cagcctggcc aacataacaa gacctcatct cttcaaaaaa tttaaaaacc
 5281 agccgggcat ggtggtgcac acctgtagtc ccagctactt gggaggctga ggtgggagga
 5341 tcacttgagt ccagcagttc aaggctgcag tgagctatgt ttgcaccacc aactccagc
 5401 ctgagtcaca gaacaagacc tcactcttaa aaaacaaaca aaaaccaaat ccacatatcc
 15 5461 taaaaaatgc tccttttcag cattctcttc tctatggaca aagggtgga tgctttaaga
 5521 accaaatctt aggctgggca cgggtggctca cgcctctaata cctagcactt tgagaggcca
 5581 aggcgggcag attgcctgag cacaggagtt cgagaccagc ctggccaaca tggtgaaacc
 5641 ctgtgtctgt caaaaataca aaaaattagc caggtgtgtt ggcgcggtgcc tataatccca
 5701 gctgctcggg aggatgaggt tcaaagaatc acttgaaccc gggaggcaga ggctgcagtg
 20 5761 agctgagatc atgccactgc actccagcct gggtgacaga gcaagacttt gtctccaaaa
 5821 aaaggaacta gacgggttca tttaaacccc tgactgcagc cctttgacat acatccaatt
 5881 gaggactggg gactccggga aacatctaaa aggccttaaaa actttgtcta acttcagccg
 5941 ggcattggtg ctacacactg taatcccagc actttgggag gctgaggcag gtggatcaca
 6001 aggtcaggag tttgagacga gcctgaccaa catggtgaaa ccccgctctc actaaaaata
 25 6061 caaaaattag ccaggcatgg tggcaggcgc ctgtaatccc agctattcgg gaggctgagg
 6121 caggagaatt gcttgaaccc cggagacaga ggttgcagcg agccgagatc gcgccactgc
 6181 actccagcct ggcaatagag tgagactcca tctcaaaaca acaacaaca caacaacaac
 6241 aacaaaatcg tctaacttcc tgatcttcc gatcattgat tttcccatag gtatgatcag
 6301 caacttgctt gggaaaggag ccgtggacca gctgacacgg ctgggtgctgg tgaatgcct
 30 6361 ctacttcaac ggccagtgga agactccctt ccccgactcc agcaccacc gccgcctctt
 6421 ccacaaatca gacggcagca ctgtctctgt gcccatgatg gctcagacca acaagttcaa
 6481 ctatagtaag tccaagagcc ccttccccac agcccacagc aactgcatct cattcctggg
 6541 gtctcccaag gaatacccaa aatgtcacc tctgaggag gaagaccaca gggaatgctc
 6601 ccctttaagg gaggagagac cctagaatat actccagctt tgacaaagat ttcccaagca
 35 6661 ggagacatca ggataatggg aacagaagac agggaggttta tcccatgaag gatgaagaag
 6721 ctgaaatcca gagattccct cagggccaca tttgtccacc tgactccagg gtctcatctt
 6781 cgtgtgttgc tagtgtgatt acctggggat gagaaatcct gctgggggag ttgagggtta
 6841 gaggatgagg actccaggtg ctgtggctca cgcctgtaat cccagcactt tgggaggcca
 6901 aggcaggtgg atcaggagtt tgaggtcagg agtttgagac cagcctggcc aacatggtga
 40 6961 aaccctgtct ctactaaaaa tgcaaaaatt agccaggtgt ggtggcaggc gcctgtaatc

7021 ccagctactc gggaggctga ggcaggagaa tcacttgagc ccgggagggtg gaggttgacg
 7081 tgagccgaac gaaattgagc cacttcaccc cagcctgggc aaaagagtga aattccattc
 7141 aaaaaaaaaa aaaaaaaaaa aaggatgagg actgggatga actggtggct ggggtgtggg
 7201 aaaatggaag tgaaggaagg caaaaagaga cagagaaggc ctggcgcggc gactcacgcc
 5 7261 tataatccca gcactttggg aggctgagaa gggggattgc ttgaggccag aagttgaata
 7321 ccagtctggg cagcatagca agaccctgcc tctacaaaaa aaaaattttt ttttaattagc
 7381 caggcttggt gacatgcac tgtagtctac tcaagaagct gaggtgaggc caggcaggggt
 7441 ggctcacgcc tgtattccca gcactttggg aggtcaaggc ggggtggatga cctgaggtca
 7501 ggagttcaag accagcctgg ccaacatggt gaaaccccat ctgtataaaa atacaaaaat
 10 7561 tagctgggca tgatagcagg tgccctgtaat tccagctact caggaggctg aggtgggaga
 7621 atctattgaa cccgggaggg ggaggttgca gtgagccgag atcatgccat tgactccag
 7681 cctgggagac agagtgagac tccttctcaa aacaaacaaa caaacaaaca aacaaaatac
 7741 agaagctgag gcgggaggaa catttgaaacc ggattcggag gctgcagtga gctatgattg
 7801 caccactgag ctccagtctg tgtgacagt agaccctgtc tcttacacac acacacacac
 15 7861 acacacacac acatgcacac acacagagag agagaaatta gaagatactg aattggcaga
 7921 agagaaggga aatagaaatt aaaatactga ataggggagc agtgaacagg ggatacccaa
 7981 aagccaagag cgagagagag cctggcttcc agaaatagtg gagaagccag gagaactagg
 8041 tgaaaaccca gtgctgggtt gccatcagcg agagctggag ccatttccaa cgaaccatct
 8101 tgtcgtcttc acagctgagt tcaccacgcc cgatggccat tactacgaca tcctggaact
 20 8161 gccctaccac ggggacaccc tcagcatggt cattgctgcc ccttatgaaa aagaggtgcc
 8221 tctctctgcc ctaccaaaca ttctgagtgc ccagctcatc agccactgga aaggcaacat
 8281 gaccaggctg ccccgccctc ttggttctgcc caagtaagcc accccgctat ctccccgacc
 8341 taccaacccc tctctcctgg ctccctaaag tcaccgcccc caggttgaat ttcccagatc
 8401 tgtgatgctt gcaggacatg catgtgtggg aggtgatgg gaaactgtgg cctgggtttg
 25 8461 attatgagtc ttgcaatcat cctccccct gtttctgctg gagggcaggg gacagctctt
 8521 cctgaccaca cccccacatt gactatcccc agaataccca gcaaaagccc ccaaaaggag
 8581 agtcagagaa atgagggagg tgggggcccc atcagtcac atctacttag ggtcgccccca
 8641 tcagcacttc catccccaac cctttcaagt caacatccaa acaaaagaaa tcacttccaa
 8701 ggacggagca gctcaaagcg cagcttctag ctgggggttc aagaaagcag atttttcgaa
 30 8761 atccttctgc agaaggaagc aaagagattt tttgaaatct ttctgcagaa ggagaaggct
 8821 ggagctgggg aactccagaa ttataggga gacctccacc acgctcatcc caaatttccg
 8881 gatgctataa tgccaggctt ggggaaagag gagaatttag ttggttagct ggtgcgtgct
 8941 ctacttgca tcctctctct tcctctttt ttttttctc ctctctctct ggctcataaa
 9001 aatggaggta attagttgtg ccctggtgag aagcagagag tgcacaaagg cccctgctt
 35 9061 gagtctctt cagggttagc tctcagaaac acaatctgca gaacagattt ttgttccaac
 9121 atccttgacg gagaatttgc ccttagcttc cccacccca gccaggctga ataaaattat
 9181 gotgaaacta ctgtcttatt tgaggaaagt aattagtcac aggtgggagg ggggtgggag
 9241 attgcagaag aatgttcag aatattagga ttttcagctc taagggggga ctttgtaaac
 9301 agctttagaa gaagaaccag gccggctggg tgtgggtggct catgcctgta atctcagcat
 40 9361 ttggggaggc caaggcgggc ggatcacttg aggtcaggag tttgagacca gcctggccaa

9421 catggtgaaa ccctgtctct attaaaaata caaaaattag ccagcgtg tagcgagcgc
 9481 ctatgatccc agctactccg gaggctgagg ccagagaatc acatgaacct gggaggtgga
 9541 ggctgcagtg agccgagatc acgccactgc actccagcct gggggacaga gcaagaatct
 9601 gtttcaaaaa aaaaaaaga aaaataggaa ggaaggaagg aaaggaaagg aaagaagaga
 5 9661 gagagaaaga aagagagaga gagagagaga aagaaagaaa gaaagaaaga aagaaagaaa
 9721 gaaagaaaga aagaaagaaa gaaaaagaaa ggaaagaaag aacgaacgaa ccaggcctcc
 9781 ctctccaacc ttcacctccg tccctattct ggccacttga ttcgggggac acctggtagg
 9841 ggatggggaa aggtgggagc tgccagccag aggggacccc ggcttgagca gcctcttgct
 9901 gctatctgca ggttctccct ggagactgaa gtcgacctca ggaagcccct agagaacctg
 10 9961 ggaatgaccg acatgttcag acagtttcag gctgacttca cgagtcttcc aggttaagaag
 10021 actttccttt gcattttctc accccagtgg actgcggggg cccctaagag gaaaaaggaa
 10081 cctctccttg agagcggcag ctgatctaat cctgtatcca catctgtttc agaccaagag
 10141 cctctccacg tcgcgaggc gctgcagaaa gtgaagatcg aggtgaacga gagtggcacg
 10201 gtggcctcct catccacagg tgagtctggc tcagggtagg ctccacgggt gtcgcctcca
 15 10261 tcgcccttca ggataactgg tcccagacc cggaaggac ccgcagccc tctcggcaca
 10321 gagcagctct gtctgtgtc agccatcacc cactccccac ctgtttctca gcctggaaaa
 10381 cgggcttggg accatggaac cctgtttcct cgcctgatgg ctccctaagtt cctgactgt
 10441 gaaaaggcct cctaaagaaa aacccaagtt gttcccacag tgggaagtaa acttaagaaa
 10501 catgcttata aggctgggca tgggtggctcc cacctgtaat ccagcgctt tgggggacca
 20 10561 aggcaggtgg atcacttgag gtcaggaatt cgagaccagc ctgggcaaca tggcaaaacc
 10621 ctatctctac taaaaataca aaaattaggc aggcgtggtg gcatgtgct gtagtcccag
 10681 ctacttggga ggctgaggca ggagaatcac ttgaatccag gaggcagagg ttgcagttag
 10741 ccgagatcac gctgctgcac tccagcctgg gcaatagagc atgactctga agaaaagaaa
 10801 gaaagaaaga gagagagaga gaaaagaaag aaagaaagaa agaaagaaag aaagaaagaa
 25 10861 agaaagaaag aaagaaagaa agagaaagaa agagaagaaa agaaaagaaa gagcttatca
 10921 ataagccctt aaaggattta gataaatgtg tgtaaggga gagctgatcc attgctacca
 10981 agtccttgga ggaaaccagg tctcagagga tgtccctaaa cttttaaggt tcatattcag
 11041 gaaaacaaac aacttcacgc tgggcttagt ggctcacacc tgtaatccca gcactttggg
 11101 aggccgaggc aggaggatcg cttgagccca ggaatttgag accagcctgg gcaatataat
 30 11161 gagactgtgc tctacaaaaa ttagaaaaaa attagccagg catggtggca tgcacctgta
 11221 gccccagtta cttgggagac tgaggtggga ggatcacttg agcccatgag ttcaaggctg
 11281 cagttagcca tgaagggtgc actgcactcc cgcctgggag acagagggag accctgtctc
 11341 taagaaaaac ggcgggggtg ggggtggtgc cagtgccagc atccctctgt tctaagacat
 11401 tgtcccttct cttgcagctg tcatagtctc agcccgcag gccccgagg agatcatcat
 35 11461 ggacagaccc ttctctttg tgggtccgca caaccccaca ggtgagcctg gaacccatca
 11521 cgttccacat cctcccaccc attttttctc tcaggaaacta gtcccagacag atgcagacat
 11581 ccctctatcc ctgagagggc tctgggcagg gaaccataac cctaccctgc ttctgtccc
 11641 aagaggaggc taccttctat caccacaga cagtgccggg tcccgcctct gtgactcagg
 11701 cagctgcgac tccagacagc tcaactcatct gcctagatct cagtccttcc acccacatcc
 40 11761 agcctgatga gctgtccac tcttctgtct tctcaacccc catggttctt ccacctcag

11821 gaacagtcct tttcatgggc caagtgatgg aaccctgacc ctggggaaag acgccttcat
11881 ctgggacaaa actggagatg catcgggaaa gaagaaactc cgaagaaaag aatttttagtg
11941 ttaatgactc tttctgaagg aagagaagac atttgccttt tgttaaaaga tggtaaacca
12001 gatctgtctc caagaccttg gcctctcctt ggaggacctt taggtcaaac tccctagtct
5 12061 ccacctgaga ccctgggaga gaagtttgaa gcacaactcc cttaaggtct ccaaaccaga
12121 cggtgacgcc tgcgggacca tctggggcac ctgcttccac ccgtctctct gccactcgg
12181 gtctgcagac ctggttccca ctgaggccct ttgcaggatg gaactacggg gcttacagga
12241 gcttttgtgt gcctggtaga aactattttct gttccagtca cattgccatc actottgtac
12301 tgcctgccac cgcggaggag gctggtgaca ggccaaaggc cagtggaaga aacacccttt
10 12361 catctcagag tccactgtgg cactggccac ccctcccag tacaggggtg ctgcagggtg
12421 cagagtgaat gtcccccac atgtggcca actctcctgg cctggccatc tccctccca
12481 gaaacagtgt gcatgggtta ttttgagtg taggtgactt gtttactcat tgaagcagat
12541 ttctgcttcc ttttattttt ataggaatag aggaagaaatg gtcagatgcg tgcccagctc
12601 ttaccccccc aatctcttgg tggggagggg tgtacctaaa tatttatcat atccttgccc
15 12661 ttgagtgtt gttagagaga aagagaacta ctaaggaaaa taatattatt taaactcgct
12721 cctagtgttt ctttgtggtc tgtgtcaccg tatctcagga agtccagcca cttgactggc
12781 acacacccct ccggacatcc agcgtgacgg agcccacact gccaccttgt ggccgcctga
12841 gacctcgcg cccccgcgc cccccgcgc cctctttttc cccttgatgg aaattgacca
12901 tacaatttca tcctccttca ggggatcaaa aggacggagt ggggggacag agactcagat
20 12961 gaggacagag tggtttccaa tgtgttcaat agatttagga gcagaaatgc aaggggctgc
13021 atgacctacc aggacagaac tttcccaat tacaggggtga ctcacagccg cattggtgac
13081 tcaactcaat gtgtcatttc cggctgctgt gtgtgagcag tggacacgtg aggggggggg
13141 tgggtgagag agacaggcag ctcgattca actaccttag ataatatattc tgaaaaccta
13201 ccagccagag ggtagggcac aaagatggat gtaatgcact ttgggaggcc aaggcgggag
25 13261 gattgcttga gcccaggagt tcaagaccag cctgggcaac ataccaagac cccgtctct
13321 ttaaaaatat atatatttta aatatactta aatatatatt tctaatatct ttaaatatat
13381 atatatattt taaagaccaa tttatgggag aattgcacac agatgtgaaa tgaatgtaat
13441 ctaatagaag cctaatacag ccacatgtt ctccactgaa aaatcctctt tctttggggt
13501 ttttctttct ttcttttttg attttgact ggacgggtgac gtcagccatg tacaggatcc
30 13561 acaggggtgg tgtcaaatgc tattgaaatt gtgttgaatt gtatgctttt tcacttttga
13621 taaataaaca tgtaaaaatg tttcaaaaaa ataataaaat aaataaatac gaagaatatg
13681 tcaggacagt cactgccttc accttctcca tttcacaccg gtgttacaag aatcagaag
13741 cctaggccag gtgtggtggt tcatgcctgt aatcccagca ctttgggaag ccgaggtggg
13801 tggatcacct aaggtcagga gtttgagacc agcctggaca acatggtgaa acccgtctc
35 13861 tactaaaaat acaaaaatta gccgggctg gtggctggcg cctgtaatcc cagctactcg
13921 ggaggctgag gcaggagaat cacttgaagc caggaggcag aggttgcagt gagctgagat
13981 tgcaccactg aactccaggc tgggtggcag agcgagactc cctctcaaaa aacaacaact
14041 acaagacaa caacaaacc agaatacaaa tcctgttggg ccatagacct catgggtgga
14101 agagaccttc ctacatccag gttggcccaa catgggggag tccatgaaat ggtcacctca
40 14161 gctctgccac aagcccaag gataagttga ttctgcccct gggaatcatc ctcaaaaagg

14221 aaaaaaatgt . tcccctgcc a taaactttcc acttatgcag atgggcctgc tcgtaagtca
14281 ctgtcaactgt gggttcccaa ctctgttcat gacacttcc tccagcacca aatgcttccc
14341 acccctctac tcccactccc cattcttcaa acccagctca agttccagtt . cctccaccta
14401 ggacttccca tggatccagc caatatcact ctcaggtcg ggcagtggc ccacgcctgt
5 14461 aatctcagca ctttgggagg ccggggcagg aagattgctt gaggccagga gtttcagacc
14521 agcctggaca acatagtgag actc

TABLE 1B

The sequences shown in TABLE 1B, are sequence fragments taken from the PAI-1 sequence shown in TABLE 1A above. Furthermore, SEQ ID NO.: 2 corresponds to positions 12571-12590 in TABLE 1A above. The nucleotide t, at position 10 in SEQ ID NO.: 2 corresponds to the nucleotide found at position 12580 of SEQ ID NO.: 1. In all of the sequences found in TABLE 2B below the polymorphism represented by the **bold** and underlined **t** may be substituted by a **g**. In SEQ ID NOS.: 3-11 in TABLE 2B, the **bold** and underlined **t** corresponds to the nucleotide found at position 12580 of SEQ ID NO.: 1. Due to the potential variability in PAI-1 sequence, the sequence motifs below may be useful in identifying PAI-1 sequences from a patient for use in genotype determination. For Example, patient sequences that form alignments with the below motifs (SEQ ID NO.: 3-11) may indicate that the patient sequence is a PAI-1 sequence and that nucleotide found at the position occupied by the **bold** and underlined **t** corresponds to the polymorphism at position 12580 of SEQ ID NO.: 1 and is therefore suitable for genotype determination. A similar strategy may be applied to the other polymorphism sites identified herein.

SEQ ID. NO.	SEQUENCE
SEQ ID. NO. 2	aggaagaaa <u>t</u> gtcagatgcg
SEQ ID. NO. 3	aggaatagag gaagaaa <u>t</u> gt
SEQ ID. NO. 4	<u>t</u> gtcagatgc gtgccagct
SEQ ID. NO. 5	aggaagaaa <u>t</u> gtcagatgcg tgcccagctc
SEQ ID. NO. 6	ataggaatag aggaagaaa <u>t</u>
SEQ ID. NO. 7	ttttatTTTT ataggaatag aggaagaaa <u>t</u>
SEQ ID. NO. 8	ttctgcttcc ttttatTTTT ataggaatag aggaagaaa <u>t</u>
SEQ ID. NO. 9	ttctgcttcc ttttatTTTT ataggaatag aggaagaaa <u>t</u> gtcagatgcg tgcccagctc
SEQ ID. NO. 10	gaagaaa <u>t</u> gt cagatg
SEQ ID. NO. 11	aagaaa <u>t</u> gtc ag

An "allele" is defined as any one or more alternative forms of a given gene. In a diploid cell or organism the members of an allelic pair (i.e. the two alleles of a given gene) occupy corresponding positions (loci) on a pair of homologous chromosomes and if these alleles are genetically identical the cell or organism is said to be "homozygous", but if genetically different the cell or organism is said to be "heterozygous" with respect to the particular gene.

A "gene" is an ordered sequence of nucleotides located in a particular position on a particular chromosome that encodes a specific functional product and may include untranslated and untranscribed sequences in proximity to the coding regions. Such non-coding sequences may contain regulatory sequences needed for transcription and translation of the sequence or introns etc.

A "genotype" is defined as the genetic constitution of an organism, usually in respect to one gene or a few genes or a region of a gene relevant to a particular context (i.e. the genetic loci responsible for a particular phenotype).

A "phenotype" is defined as the observable characters of an organism.

A "single nucleotide polymorphism" (SNP) occurs at a polymorphic site occupied by a single nucleotide, which is the site of variation between allelic sequences. The site is usually preceded by and followed by highly conserved sequences of the allele (e.g., sequences that vary in less than 1/100 or 1/1000 members of the populations). A single nucleotide polymorphism usually arises due to substitution of one nucleotide for another at the polymorphic site. A "transition" is the replacement of one purine by another purine or one pyrimidine by another pyrimidine. A "transversion" is the replacement of a purine by a pyrimidine or vice versa. Single nucleotide polymorphisms can also arise from a deletion (represented by "-" or "del") of a nucleotide or an insertion (represented by "+" or "ins") of a nucleotide relative to a reference allele. Furthermore, it would be appreciated by a person of skill in the art, that an insertion or deletion within a given sequence could alter the relative position and therefore the position number of another polymorphism within the sequence.

A "systemic inflammatory response syndrome" (SIRS) is defined as including both septic (i.e. sepsis or septic shock) and non-septic systemic inflammatory response (i.e. post operative).

“SIRS” is further defined according to ACCP (American College of Chest Physicians) guidelines as the presence of two or more of A) temperature $> 38^{\circ}\text{C}$ or $< 36^{\circ}\text{C}$, B) heart rate > 90 beats per minute, C) respiratory rate > 20 breaths per minute, and D) white blood cell count $> 12,000$ per mm^3 or $< 4,000$ mm^3 . In the following description, the presence of two, three, or four of the “SIRS” criteria were scored each day over the 28 day observation period.

“Sepsis” is defined as the presence of at least two “SIRS” criteria and known or suspected source of infection. Septic shock was defined as sepsis plus one new organ failure by Brussels criteria plus need for vasopressor medication.

Patient outcome or prognosis as used herein refers the ability of a patient to recover from an inflammatory condition. An inflammatory condition, may be selected from the group consisting of: sepsis, septicemia, pneumonia, septic shock, systemic inflammatory response syndrome (SIRS), Acute Respiratory Distress Syndrome (ARDS), acute lung injury, infection, pancreatitis, bacteremia, peritonitis, abdominal abscess, inflammation due to trauma, inflammation due to surgery, chronic inflammatory disease, ischemia, ischemia-reperfusion injury of an organ or tissue, tissue damage due to disease, tissue damage due to chemotherapy or radiotherapy, and reactions to ingested, inhaled, infused, injected, or delivered substances, glomerulonephritis, bowel infection, opportunistic infections, and for patients undergoing major surgery or dialysis, patients who are immunocompromised, patients on immunosuppressive agents, patients with HIV/AIDS, patients with suspected endocarditis, patients with fever, patients with fever of unknown origin, patients with cystic fibrosis, patients with diabetes mellitus, patients with chronic renal failure, patients with bronchiectasis, patients with chronic obstructive lung disease, chronic bronchitis, emphysema, or asthma, patients with febrile neutropenia, patients with meningitis, patients with septic arthritis, patients with urinary tract infection, patients with necrotizing fasciitis, patients with other suspected Group A streptococcus infection, patients who have had a splenectomy, patients with recurrent or suspected enterococcus infection, other medical and surgical conditions associated with increased risk of infection, Gram positive sepsis, Gram negative sepsis, culture negative sepsis, fungal sepsis, meningococcemia, post-pump syndrome, cardiac stun syndrome, myocardial infarction, stroke, congestive heart failure, hepatitis, epiglottitis, E. coli 0157:H7, malaria, gas gangrene, toxic shock syndrome, mycobacterial tuberculosis, Pneumocystic carinii, pneumonia, Leishmaniasis, hemolytic uremic syndrome/thrombotic thrombocytopenic purpura, Dengue hemorrhagic fever, pelvic

inflammatory disease, Legionella, Lyme disease, Influenza A, Epstein-Barr virus, encephalitis, inflammatory diseases and autoimmunity including Rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, inflammatory bowel disease, idiopathic pulmonary fibrosis, sarcoidosis, hypersensitivity pneumonitis, systemic vasculitis, Wegener's granulomatosis, transplants including heart, liver, lung kidney bone marrow, graft-versus-host disease, transplant rejection, sickle cell anemia, nephrotic syndrome, toxicity of agents such as OKT3, cytokine therapy, and cirrhosis.

Assessing patient outcome or prognosis may be accomplished by various methods. For Example, an "APACHE II" score is defined as Acute Physiology And Chronic Health Evaluation and herein was calculated on a daily basis from raw clinical and laboratory variables. (VINCENT *et al. Scoring systems for assessing organ dysfunction and survival*. Critical Care Clinics. 16:353-366, 2000) summarize APACHE score as follows "First developed in 1981 by KNAUS WA *et al. (Chest (1991) 100(6):1619-36)*, the APACHE score has become the most commonly used survival prediction model in ICUs worldwide. The APACHE II score, a revised and simplified version of the original prototype, uses a point score based on initial values of 12 routine physiologic measures, age, and previous health status to provide a general measure of severity of disease. The values recorded are the worst values taken during the patient's first 24 hours in the ICU. The score is applied to one of 34 admission diagnoses to estimate a disease-specific probability of mortality (APACHE II predicted risk of death). The maximum possible APACHE II score is 71, and high scores have been well correlated with mortality. The APACHE II score has been widely used to stratify and compare various groups of critically ill patients, including patients with sepsis, by severity of illness on entry into clinical trials."

A "Brussels score" score is a method for evaluating organ dysfunction as compared to a baseline. If the Brussels score is 0 (ie. moderate, severe, or extreme), then organ failure was recorded as present on that particular day (see TABLE 1C below). In the following description, to correct for deaths during the observation period, days alive and free of organ failure (DAF) were calculated as previously described. For example, acute lung injury was calculated as follows. Acute lung injury is defined as present when a patient meets all of these four criteria. 1) Need for mechanical ventilation, 2) Bilateral pulmonary infiltrates on chest X-ray consistent with acute lung injury, 3) $\text{PaO}_2/\text{FiO}_2$ ratio is less than 300, 4) No clinical evidence of congestive heart failure or if a pulmonary artery catheter is in place for clinical purposes, a pulmonary capillary wedge pressure

less than 18 mm Hg (1). The severity of acute lung injury is assessed by measuring days alive and free of acute lung injury over a 28 day observation period. Acute lung injury is recorded as present on each day that the person has moderate, severe or extreme dysfunction as defined in the Brussels score. Days alive and free of acute lung injury is calculated as the number of days after onset of acute lung injury that a patient is alive and free of acute lung injury over a defined observation period (28 days). Thus, a lower score for days alive and free of acute lung injury indicates more severe acute lung injury. The reason that days alive and free of acute lung injury is preferable to simply presence or absence of acute lung injury, is that acute lung injury has a high acute mortality and early death (within 28 days) precludes calculation of the presence or absence of acute lung injury in dead patients. The cardiovascular, renal, neurologic, hepatic and coagulation dysfunction were similarly defined as present on each day that the person had moderate, severe or extreme dysfunction as defined by the Brussels score. Days alive and free of steroids are days that a person is alive and is not being treated with exogenous corticosteroids (e.g. hydrocortisone, prednisone, methylprednisolone). Days alive and free of pressors are days that a person is alive and not being treated with intravenous vasopressors (e.g. dopamine, norepinephrine, epinephrine, phenylephrine). Days alive and free of an International Normalized Ratio (INR) > 1.5 are days that a person is alive and does not have an INR > 1.5.

TABLE 1C

Brussels Organ Dysfunction Scoring System

ORGANS	Free of Organ Dysfunction		Clinically Significant Organ Dysfunction		
	Normal	Mild	Moderate	Severe	Extreme
DAF ORGAN DYSFUNCTION SCORE	1		0		
<u>Cardiovascular</u> Systolic BP (mmHg)	>90	≤90 Responsive to fluid	≤90 Unresponsive to fluid	≤90 plus pH ≤7.3	≤90 plus pH ≤7.2
<u>Pulmonary</u> P _a O ₂ /F _i O ₂ (mmHg)	>400	400-301	300-201 Acute lung injury	200-101 ARDS	≤100 Severe ARDS
<u>Renal</u> Creatinine (mg/dL)	<1.5	1.5-1.9	2.0-3.4	3.5-4.9	≥5.0
<u>Hepatic</u> Bilirubin (mg/dL)	<1.2	1.2-1.9	2.0-5.9	6.0-11.9	≥12

<u>Hematologic</u> Platelets ($\times 10^5/\text{mm}^3$)	>120	120-81	80-51	50-21	≤ 20
<u>Neurologic</u> (Glasgow Score)	15	14-13	12-10	9-6	≤ 5
Round Table Conference on Clinical Trials for the Treatment of Sepsis Brussels, March 12-14, 1994.					

Analysis of variance (ANOVA) is a standard statistical approach to test for statistically significant differences between sets of measurements.

- 5 The Fisher exact test is a standard statistical approach to test for statistically significant differences between rates and proportions of characteristics measured in different groups.

2. General Methods

One aspect of the invention may involve the identification of patients or the selection of patients
 10 that are either at risk of developing an inflammatory condition or the identification of patients who already have an inflammatory condition. For example, patients who have undergone major surgery or scheduled for or contemplating major surgery may be considered as being at risk of developing an inflammatory condition. Furthermore, patients may be determined as having an inflammatory condition using diagnostic methods and clinical evaluations known in the medical
 15 arts. An inflammatory condition, may be selected from the group consisting of: sepsis, septicemia, pneumonia, septic shock, systemic inflammatory response syndrome (SIRS), Acute Respiratory Distress Syndrome (ARDS), acute lung injury, infection, pancreatitis, bacteremia, peritonitis, abdominal abscess, inflammation due to trauma, inflammation due to surgery, chronic inflammatory disease, ischemia, ischemia-reperfusion injury of an organ or tissue, tissue damage
 20 due to disease, tissue damage due to chemotherapy or radiotherapy, and reactions to ingested, inhaled, infused, injected, or delivered substances, glomerulonephritis, bowel infection, opportunistic infections, and for patients undergoing major surgery or dialysis, patients who are immunocompromised, patients on immunosuppressive agents, patients with HIV/AIDS, patients with suspected endocarditis, patients with fever, patients with fever of unknown origin, patients
 25 with cystic fibrosis, patients with diabetes mellitus, patients with chronic renal failure, patients with bronchiectasis, patients with chronic obstructive lung disease, chronic bronchitis, emphysema, or asthma, patients with febrile neutropenia, patients with meningitis, patients with septic arthritis, patients with urinary tract infection, patients with necrotizing fasciitis, patients

with other suspected Group A streptococcus infection, patients who have had a splenectomy, patients with recurrent or suspected enterococcus infection, other medical and surgical conditions associated with increased risk of infection, Gram positive sepsis, Gram negative sepsis, culture negative sepsis, fungal sepsis, meningococcemia, post-pump syndrome, cardiac stun syndrome, myocardial infarction, stroke, congestive heart failure, hepatitis, epiglottitis, *E. coli* 0157:H7, malaria, gas gangrene, toxic shock syndrome, mycobacterial tuberculosis, *Pneumocystis carinii*, pneumonia, Leishmaniasis, hemolytic uremic syndrome/thrombotic thrombocytopenic purpura, Dengue hemorrhagic fever, pelvic inflammatory disease, *Legionella*, Lyme disease, Influenza A, Epstein-Barr virus, encephalitis, inflammatory diseases and autoimmunity including Rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, inflammatory bowel disease, idiopathic pulmonary fibrosis, sarcoidosis, hypersensitivity pneumonitis, systemic vasculitis, Wegener's granulomatosis, transplants including heart, liver, lung kidney bone marrow, graft-versus-host disease, transplant rejection, sickle cell anemia, nephrotic syndrome, toxicity of agents such as OKT3, cytokine therapy, and cirrhosis.

Once a patient is identified as being at risk for developing or having an inflammatory condition, then genetic sequence information may be obtained from the patient. Or alternatively genetic sequence information may already have been obtained from the patient. For example, a patient may have already provided a biological sample for other purposes or may have even had their genetic sequence determined in whole or in part and stored for future use. Genetic sequence information may be obtained in numerous different ways and may involve the collection of a biological sample that contains genetic material. Particularly, genetic material, containing the sequence or sequences of interest. Many methods are known in the art for collecting bodily samples and extracting genetic material from those samples. Genetic material can be extracted from blood, tissue and hair and other samples. There are many known methods for the separate isolation of DNA and RNA from biological material. Typically, DNA may be isolated from a biological sample when first the sample is lysed and then the DNA is isolated from the lysate according to any one of a variety of multi-step protocols, which can take varying lengths of time. DNA isolation methods may involve the use of phenol (SAMBROOK, J. *et al.*, "Molecular Cloning", Vol. 2, pp. 9.14-9.23, Cold Spring Harbor Laboratory Press (1989) and AUSUBEL, FM. *et al.*, "Current Protocols in Molecular Biology", Vol. 1, pp. 2.2.1-2.4.5, John Wiley & Sons, Inc. (1994)). Typically, a biological sample is lysed in a detergent solution and the protein component of the lysate is digested with proteinase for 12-18 hours. Next, the lysate is extracted

with phenol to remove most of the cellular components, and the remaining aqueous phase is processed further to isolate DNA. In another method, described in VAN NESS *et al.* (U.S. Pat. # 5,130,423), non-corrosive phenol derivatives are used for the isolation of nucleic acids. The resulting preparation is a mix of RNA and DNA.

5

Other methods for DNA isolation utilize non-corrosive chaotropic agents. These methods, which are based on the use of guanidine salts, urea and sodium iodide, involve lysis of a biological sample in a chaotropic aqueous solution and subsequent precipitation of the crude DNA fraction with a lower alcohol. The final purification of the precipitated, crude DNA fraction can be achieved by any one of several methods, including column chromatography (ANALECTS, (1994) Vol 22, No. 4, Pharmacia Biotech), or exposure of the crude DNA to a polyanion-containing protein as described in KOLLER (U.S. Pat. # 5,128,247).

10

Yet another method of DNA isolation, which is described by BOTWELL DDL (Anal. Biochem. (1987) 162:463-465) involves lysing cells in 6M guanidine hydrochloride, precipitating DNA from the lysate at acid pH by adding 2.5 volumes of ethanol, and washing the DNA with ethanol.

15

Numerous other methods are known in the art to isolate both RNA and DNA, such as the one described by CHOMCZYNSKI (U.S. Pat. # 5,945,515), whereby genetic material can be extracted efficiently in as little as twenty minutes. EVANS and HUGH (U.S. Pat. # 5,989,431) describe methods for isolating DNA using a hollow membrane filter.

20

Once a patient's genetic sequence information has been obtained from the patient it may then be further analyzed to detect or determine the identity or genotype of one or more polymorphisms in the PAI-1 gene. Provided that the genetic material obtained, contains the sequence of interest. Particularly, a person may be interested in determining the PAI-1 promoter genotype of a patient of interest, where the genotype includes a nucleotide corresponding to position 12580 or SEQ ID NO.: 1 or position 8 of SEQ ID NO.: 2. The sequence of interest may also include other PAI-1 gene polymorphisms or may also contain some of the sequence surrounding the polymorphism of interest. Detection or determination of a nucleotide identity or the genotype of the single nucleotide polymorphism(s) or other polymorphism, may be accomplished by any one of a number methods or assays known in the art, including but not limited to the following:

25
30

Restriction Fragment Length Polymorphism (RFLP) strategy – An RFLP gel-based analysis can be used to distinguish between alleles at polymorphic sites within a gene. Briefly, a short segment of DNA (typically several hundred base pairs) is amplified by PCR. Where possible, a specific restriction endonuclease is chosen that cuts the short DNA segment when one variant allele is present but does not cut the short DNA segment when the other allele variant is present. After incubation of the PCR amplified DNA with this restriction endonuclease, the reaction products are then separated using gel electrophoresis. Thus, when the gel is examined the appearance of two lower molecular weight bands (lower molecular weight molecules travel farther down the gel during electrophoresis) indicates that the initial DNA sample had the allele which could be cut by the chosen restriction endonuclease. In contrast, if only one higher molecular weight band is observed (at the molecular weight of the PCR product) then the initial DNA sample had the allele variant that could not be cut by the chosen restriction endonuclease. Finally, if both the higher molecular weight band and the two lower molecular weight bands are visible then the initial DNA sample contained both alleles, and therefore the patient was heterozygous for this single nucleotide polymorphism;

Sequencing – For example the Maxam-Gilbert technique for sequencing (MAXAM AM. and GILBERT W. Proc. Natl. Acad. Sci. USA (1977) 74(4):560-564) involves the specific chemical cleavage of terminally labelled DNA. In this technique four samples of the same labeled DNA are each subjected to a different chemical reaction to effect preferential cleavage of the DNA molecule at one or two nucleotides of a specific base identity. The conditions are adjusted to obtain only partial cleavage, DNA fragments are thus generated in each sample whose lengths are dependent upon the position within the DNA base sequence of the nucleotide(s) which are subject to such cleavage. After partial cleavage is performed, each sample contains DNA fragments of different lengths, each of which ends with the same one or two of the four nucleotides. In particular, in one sample each fragment ends with a C, in another sample each fragment ends with a C or a T, in a third sample each ends with a G, and in a fourth sample each ends with an A or a G. When the products of these four reactions are resolved by size, by electrophoresis on a polyacrylamide gel, the DNA sequence can be read from the pattern of radioactive bands. This technique permits the sequencing of at least 100 bases from the point of labeling. Another method is the dideoxy method of sequencing was published by Sanger *et al.*

(SANGER *et al.* Proc. Natl. Acad. Sci. USA (1977) 74(12):5463-5467). The Sanger method relies on enzymatic activity of a DNA polymerase to synthesize sequence-dependent fragments of various lengths. The lengths of the fragments are determined by the random incorporation of dideoxynucleotide base-specific terminators. These fragments can then be separated in a gel as in the Maxam-Gilbert procedure, visualized, and the sequence determined. Numerous improvements have been made to refine the above methods and to automate the sequencing procedures. Similarly, RNA sequencing methods are also known. For example, reverse transcriptase with dideoxy-nucleotides have been used to sequence encephalomyocarditis virus RNA (ZIMMERN D. and KAESBERG P. Proc. Natl. Acad. Sci. USA (1978) 75(9):4257-4261). MILLS DR. and KRAMER FR. (Proc. Natl. Acad. Sci. USA (1979) 76(5):2232-2235) describe the use of Q.beta. replicase and the nucleotide analog inosine for sequencing RNA in a chain-termination mechanism. Direct chemical methods for sequencing RNA are also known (PEATTIE DA. Proc. Natl. Acad. Sci. USA (1979) 76(4):1760-1764). Other methods include those of DONIS-KELLER *et al.* (1977, Nucl. Acids Res. 4:2527-2538), SIMONCSITS A. *et al.* (Nature (1977) 269(5631):833-836), AXELROD VD. *et al.* (Nucl. Acids Res.(1978) 5(10):3549-3563), and KRAMER FR. and MILLS DR. (Proc. Natl. Acad. Sci. USA (1978) 75(11):5334-5338, which are incorporated herein by reference). Nucleic acid sequences can also be read by stimulating the natural fluoresce of a cleaved nucleotide with a laser while the single nucleotide is contained in a fluorescence enhancing matrix (U.S. Pat. # 5,674,743);

Hybridization methods for the identification of SNPs using hybridization techniques are described in the U.S. Pat. # 6,270,961 & 6,025,136;

A template-directed dye-terminator incorporation with fluorescent polarization-detection (TDI-FP) method is described by FREEMAN BD. *et al.* (J Mol Diagnostics (2002) 4(4):209-215) is described for large scale screening;

Oligonucleotide ligation assay (OLA) - is based on ligation of probe and detector oligonucleotides annealed to a polymerase chain reaction amplicon strand with detection by an enzyme immunoassay (VILLAHERMOSA ML. J Hum Virol (2001) 4(5):238-48;

ROMPPANEN EL. Scand J Clin Lab Invest (2001) 61(2):123-9; IANNONE MA. *et al.* Cytometry (2000) 39(2):131-40;

Ligation-Rolling Circle Amplification (L-RCA) has also been successfully used for genotyping single nucleotide polymorphisms as described in QI X. *et al.* Nucleic Acids Res (2001) 29(22):E116;

5' nuclease assay has also been successfully used for genotyping single nucleotide polymorphisms (AYDIN A. *et al.* Biotechniques (2001) (4):920-2, 924, 926-8.);

Polymerase proofreading methods are used to determine SNPs identities, as described in WO 0181631;

Detection of single base pair DNA mutations by enzyme-amplified electronic transduction is described in PATOLSKY F *et al.* Nat Biotech. (2001) 19(3):253-257; or

Allele specific PCR methods have also been successfully used for genotyping single nucleotide polymorphisms (HAWKINS JR. *et al.* Hum Mutat (2002) 19(5):543-553).

Alternatively, if a patient's sequence data is already known, then obtaining may involve retrieval of the patients nucleic acid sequence data from a database, followed by determining or detecting the identity of a nucleic acid or genotype at a polymorphism site by reading the patient's nucleic acid sequence at the polymorphic site.

Once the identity of a polymorphism(s) is determined or detected an indication may be obtained as to patient outcome or prognosis based on the genotype (the nucleotide identity at a position) of the polymorphism of interest. In the present invention, polymorphisms in the PAI-1 gene, are used to obtain a prognosis or to determine patient outcome or selection of therapies. Methods for obtaining patient outcome or prognosis or for patient screening may be useful to determine the ability of a patient to recover from an inflammatory condition. Alternatively, single polymorphism sites or combined polymorphism sites may be used as an indication of a patient's ability to recover from an inflammatory condition, if they are linked to a polymorphism determined to be indicative of a patient's ability to recover from an inflammatory condition.

Once patient outcome or a prognosis is determined, such information may be of interest to physicians and surgeons to assist in deciding between potential treatment options, to help determine the degree to which patients are monitored and the frequency with which such monitoring occurs. Ultimately, treatment decisions may be made in response to factors, both specific to the patient and based on the experience of the physician or surgeon responsible for a patient's care. Treatment options that a physician or surgeon may consider in treating a patient with an inflammatory condition may include, but are not limited to the following:

- (a) use of anti-inflammatory therapy;
- (b) use of steroids;
- (c) use of activated Protein C (drotrocogin alpha or Xigris™ from Lilly);
- (d) use of modulators of the coagulation cascade (such as various versions of heparin) use of antibody to tissue factor;
- (e) use of anti-thrombin or anti-thrombin III;
- (f) streptokinase;
- (g) use of antiplatelet agents such as clopidogrel; and
- (h) Surfactant.

Alternative treatments currently in development and potentially useful in the treatment of an inflammatory condition may include, but are not limited to the following: antibodies to tumor necrosis factor (TNF) or even antibody to endotoxin (i.e. lipopolysaccharide, LPS); tumor necrosis factor receptor (TNF); tissue factor pathway inhibitors (tifacogin™ alpha from Chiron); platelet activating factor hydrolase (PAFase™ from ICOS); antibodies to IL-6; antibodies, antagonists or inhibitors to high mobility group box 1 (HMGB-1 or HMG-1 tissue plasminogen activator; bradykinin antagonists; antibody to CD-14; interleukin-10; Recombinant soluble tumor necrosis factor receptor-immunoglobulin G1(Roche); Procysteine; Elastase Inhibitor; and human recombinant interleukin 1 receptor antagonist (IL-1 RA).

Alternatively, similar methods may be employed to identify new polymorphisms in PAI-1 sequence that correlate with patient outcome or prognosis.

As described above genetic sequence information or genotype information may be obtained from a patient wherein the sequence information contains one or more single nucleotide polymorphism sites in the PAI-1 gene. Also, as previously described the sequence identity of one or more single nucleotide polymorphisms in the PAI-1 gene of one or more patients may then be detected or determined. Furthermore, patient outcome or prognosis may be assessed as described above, for example the APACHE II scoring system or the Brussels score may be used to assess patient outcome or prognosis by comparing patient scores before and after treatment. Once patient outcome or prognosis has been assessed, patient outcome or prognosis may be correlated with the sequence identity of a single nucleotide polymorphism(s). The correlation of patient outcome or prognosis may further include statistical analysis of patient outcome scores and polymorphism(s) for a number of patients.

Clinical Phenotype

The primary outcome variable was survival to hospital discharge. Secondary outcome variables were days alive and free of cardiovascular, respiratory, renal, hepatic, hematologic, and neurologic organ system failure as well as days alive and free of SIRS (Systemic Inflammatory Response Syndrome), occurrence of sepsis, and occurrence of septic shock. SIRS was considered present when patients met at least two of four SIRS criteria. The SIRS criteria were 1) fever ($>38^{\circ}\text{C}$) or hypothermia ($<35.5^{\circ}\text{C}$), 2) tachycardia (>100 beats/min in the absence of beta blockers, 3) tachypnea (>20 breaths/min) or need for mechanical ventilation, and 4) leukocytosis (total leukocyte count $> 11,000/\mu\text{L}$) (Anonymous. *Critical Care Medicine* (1992) 20(6):864-74). Patients were included in this cohort on the calendar day on which the SIRS criteria were met.

A patients' baseline demographics that were recorded included age, gender, whether medical or surgical diagnosis for admission (according to APACHE III diagnostic codes (KNAUS WA et al. *Chest* (1991) 100(6):1619-36)), and admission APACHE II score. The following additional data were recorded for each 24 hour period (8 am to 8 am) for 28 days to evaluate organ dysfunction, SIRS, sepsis, and septic shock.

Clinically significant organ dysfunction for each organ system was defined as present during a 24 hour period if there was evidence of at least moderate organ dysfunction using the Brussels

criteria (TABLE 1C) (RUSSELL JA et al. *Critical Care Medicine* (2000) 28(10):3405-11). Because data were not always available during each 24 hour period for each organ dysfunction variable, we used the "carry forward" assumption as defined previously (Anonymous. *New England Journal of Medicine* (2000) 342(18):1301-8). Briefly, for any 24 hour period in which there was no measurement of a variable, we carried forward the "present" or "absent" criteria from the previous 24 hour period. If any variable was never measured, it was assumed to be normal.

To further evaluate cardiovascular, respiratory, and renal function we also recorded, during each 24 hour period, vasopressor support, mechanical ventilation, and renal support, respectively. Vasopressor use was defined as dopamine > 5 µg/kg/min or any dose of norepinephrine, epinephrine, vasopressin, or phenylephrine. Mechanical ventilation was defined as need for intubation and positive airway pressure (i.e. T- piece and mask ventilation were not considered ventilation). Renal support was defined as hemodialysis, peritoneal dialysis, or any continuous renal support mode (e.g. continuous veno-venous hemodialysis).

To assess duration of organ dysfunction and to correct organ dysfunction scoring for deaths in the 28-day observation period, calculations were made of days alive and free of organ dysfunction (DAF) as previously reported (BERNARD GR et al. *New England Journal of Medicine* (1997) 336(13):912-8). Briefly, during each 24-hour period for each variable, DAF was scored as 1 if the patient was alive and free of organ dysfunction (normal or mild organ dysfunction, Table 1). DAF was scored as 0 if the patient had organ dysfunction (moderate, severe, or extreme) or was not alive during that 24-hour period. Each of the 28 days after ICU admission was scored in each patient in this fashion. Thus, the lowest score possible for each variable was zero and the highest score possible was 28. A low score is indicative of more organ dysfunction as there would be fewer days alive and free of organ dysfunction.

Similarly, days alive and free of SIRS (DAF SIRS) were calculated. Each of the four SIRS criteria were recorded as present or absent during each 24 hour period. Presence of SIRS during each 24 hour period was defined by having at least 2 of the 4 SIRS criteria. Sepsis was defined as present during a 24 hour period by having at least two of four SIRS criteria and having a known or suspected infection during the 24 hour period (Anonymous. *Critical Care Medicine* (1992)

20(6):864-74). Cultures that were judged to be positive due to contamination or colonization were excluded. Septic shock was defined as presence of sepsis plus presence of hypotension (systolic blood pressure < 90 mmHg or need for vasopressor agents) during the same 24 hour period.

5

Haplotypes and Selection of htSNPs

Using unphased Caucasian genotypic data (from pga.mbt.washington.edu (RIEDER MJ et al. ACCESSION AF386492, Vol. 2001: SeattleSNPs. NHLBI Program for Genomic Applications, UW-FHCRC, Seattle, WA (2001)) haplotypes were inferred using PHASE software (Figure 1).
10 MEGA 2 was then used to infer a phylogenetic tree to identify major haplotype clades (Figure 2). Haplotypes were sorted according to the phylogenetic tree and haplotype structure was inspected to choose haplotype tag SNPs (htSNPs). It was found that 4 htSNPs uniquely identified the major haplotype clades (Figures 1 and 2). The 4G/5G polymorphism, which corresponds to -675 in DAWSON et al. (1993) and position 837 of SEQ ID NO:1 (NCBI ID: rs1799889) and four
15 htSNPs: 11312 (9732) G/A, 12580 (11000) T/G; 12750 (11170) G/A; 13985 (12405) C/T (bracketed positions numbered according to the start (1581) of the PAI-1 protein coding region - unbracketed positions numbered according of the sequence represented by accession number AF386492) were then genotyped in the patient cohort. Finally, PHASE was used to infer haplotypes from the genotyped htSNPS in each patient in the critically ill cohort with SIRS.

20

Genotyping

DNA was extracted from peripheral blood samples using a QIAamp DNA Blood Maxi Kit (Qiagen). The 4G/5G polymorphism was genotyped using a restriction fragment length strategy. The htSNPs were genotyped using Masscode tagging (Qiagen Genomics, Inc) (KOKORIS M et al
25 *Molecular Diagnosis* (2000) 5(4):329-40). The genotyping success rate was >99% for all SNPs.

3. EXAMPLES

Patient Cohort for Examples 1 and 2

Data were collected for 235 consecutive patients who were admitted to the Intensive Care Unit
30 (ICU) of St. Paul's Hospital. This ICU is a mixed medical – surgical ICU in a tertiary care, university-affiliated teaching hospital of the University of British Columbia. Results are reported for only the Caucasian patients (78.6% of all admissions) who were successfully genotyped (n=223) in order to decrease the potential confounding influence of population admixture

secondary to ethnic diversity, on associations between genotype and phenotype. These 235 patients make up the cohort of this study.

Example 1 - Haplotype clade deduction

5 First haplotypes of the PAI-1 gene (Figure 1) were deduced, the major haplotype clades were identified (Figures 1 and 2), and a minimum set of htSNPs defining the major haplotype clades (Figure 1) were selected. The htSNPs and the 4G/5G polymorphism were in Hardy-Weinberg equilibrium (Table 2). Furthermore, it was found that haplotypes containing the 4G allele could be grouped into the allele that contained the minor T allele of a C/T polymorphism at position
10 13985 and the allele that contained the minor G allele of a T/G polymorphism at position 12580 (Figures 1 and 2). The haplotype map in Figure 1, shows PAI-1 gene haplotypes in Caucasians. Haplotypes were inferred using PHASE from available data, and represented in the style of PATIL et al. (Science (2001) 294(5547):1669-70). Columns indicate polymorphic loci and rows indicate distinct haplotypes inferred in Caucasian individuals. Light cells indicate rare alleles, and
15 dark cells indicate common alleles. The map was generated from genotype data (available publicly at <http://pga.mbt.washington.edu/>) of 23 unrelated Caucasians from the Coriell registry of patients. Haplotypes were inferred using the statistical program PHASE 2.0 (STEPHENS M and DONNELLY P Am J Human Genet. (2003) 73(6):1162-1169). Polymorphisms that provide the same information as the 12580G allele were chosen based on this haplotype map. These
20 polymorphisms provide the same information as the 12580G allele. One method is to positively identify the 12580G allele (section A) either directly or in combination. Another method is to positively identify the 12580T allele (section B), either directly or in combination, and thereby negatively identify the 12580G allele. Haplotypes within each clade are very similar, while clades differ substantially from each other. MEGA II was used to sort haplotypes into clades
25 separated by heavier lines.

An unrooted phylogenetic tree of PAI-1 haplotype clades identified in Figure 1, is shown in Figure 2. Figure 2 was drawn using MEGA II where line length reflects relative evolutionary distance. Each clade, shown with circles, is identified by haplotype tag SNPs (htSNPs) which are
30 representative of specific clades. Recombination of the 4G allele from its ancestral clade (htSNP allele 12580 G) appears to have occurred with an evolutionarily very distant clade (htSNP allele 13985 T).

TABLE 2
Genotype and Allele Frequencies for 4G/5G and Four htSNPS in a Cohort of Critically Ill Adults who had SIRS

	Genotype Frequencies			Allele Frequencies	
	4G/4G	4G/5G	5G/5G	4G	5G
837 (4G/5G)	0.27	0.53	0.20	0.53	0.47
	GG	GA	AA	G	A
11312 G/A (9732)	0.77	0.21	0.02	0.88	0.13
	TT	TG	GG	T	G
12580 T/G (11000)	0.35	0.49	0.16	0.59	0.41
	GG	GA	AA	G	A
12750 G/A (11170)	0.63	0.34	0.03	0.80	0.20
	CC	CT	TT	C	T
13985 C/T (12405)	0.81	0.17	0.02	0.90	0.10

5

Example 2 - Haplotype Clade Patient Outcome 4G Allele v. PAI-1 Clades

As shown in Figure 3, two of the six haplotype clades, D and F, are combined to make up the 4G allele and defined by: the 13985 T and the 12580 G minor alleles respectively. It was found that patients having both of the 4G haplotype clades did not demonstrate decreased hospital survival (Figure 3). In fact, the clade containing the 13985 T allele had the highest survival of all clades - in direct contrast to reported results for the 4G allele (HERMANS (1999); MENGES (2001); and WESTENDORP (1999)). The 4G (875) allele was similarly not associated with altered survival, organ dysfunction, SIRS, sepsis, or septic shock. In contrast, those patients having 12580 G allele had markedly decreased hospital survival compared to the 13985 T allele ($p < 0.01$) and compared to all other haplotype clades ($p < 0.03$) (Figure 3).

Example 3 - Patient Outcomes for G4/G5 Genotypes

When only the 4G/5G polymorphism was examined, intermediate results reflecting the addition of the increased survival of the haplotype clade, containing 13985 T allele, to the decreased survival of the haplotype clade, containing the 12580 G allele, were found. The frequency distribution of the 4G/4G, 4G/5G and 5G/5G genotypes were 26.6%, 52.1% and 20.1% which is similar to the frequencies reported in other critically ill Caucasians (MENGES T et al. *Lancet* (2001) 357(9262):1096-7). Hospital survival rates were 54% for 4G/4G, 61% for 4G/5G, and 74% for

5G/5G groups, which were not statistically significantly different ($p = 0.09$). However, the 4G allele remained associated with significantly fewer days alive and free of respiratory dysfunction ($p < 0.05$), hepatic dysfunction ($p < 0.05$), and neurologic dysfunction ($p < 0.05$) (TABLE 3).

Cardiovascular, renal, and coagulation DAF scores were not significantly different between

4G/5G genotypes (Table 3). These results were supported by the findings of no difference in DAF vasopressors (additional measurement of cardiovascular function) but significantly fewer DAF ventilator (additional measurement of respiratory function) in patients who had the 4G allele (Table 3). Those patients who had the 4G allele were found to have a significantly increased duration of SIRS (DAF SIRS: 19.8 out of 28 for 4G/4G, 20.8 for 4G/5G, 23.4 for 5G/5G), a trend to increased occurrence of sepsis (94% for 4G/4G, 99% for 4G/5G, 74% for 5G/5G, $p=0.08$) and a significant increase in occurrence of septic shock (94% for 4G/4G, 99% for 4G/5G, 74% for 5G/5G, $p=0.04$) during the 28 day study period.

TABLE 3
Days Alive and Free of Organ Failure (out of a possible 28)
for PAI-1 4G/5G Polymorphism Genotypes in a Cohort of Critically Ill Adults who had SIRS

	4G / 4G	4G / 5G	5G / 5G	p
Cardiovascular	18.5±2.6	19.7±1.8	22.3±2.7	0.07
Vasopressors (marker of Cardiovascular dysfunction)	17.2±2.6	18.6±1.9	21.0±2.8	0.08
Respiratory	7.2±2.1	5.9±1.4	12.3±2.9	0.02
Ventilator (marker of Respiratory dysfunction)	12.7±2.8	13.9±1.9	17.0±2.9	0.04
Renal	16.5±2.7	16.3±2.0	18.0±3.3	0.52
Renal Support (marker of Renal dysfunction)	18.7±2.7	18.1±2.0	19.4±3.2	0.85
Hepatic	18.0±2.8	19.0±1.9	23.0±2.8	0.02
Hematologic				
Neurologic	18.4±2.6	19.5±1.9	22.8±2.6	0.04

Mean ± 95% Confidence Interval

Example 4 - Patient Outcomes for PAI-1 12580 Genotypes v. 4G/5G Genotypes

To assess duration of organ dysfunction and to correct organ dysfunction scoring for deaths in the 28-day observation period, days alive and free of organ dysfunction (DAF) were calculated.

During each 24-hour period for each variable, DAF was scored as "1" if the patient was alive and free of organ dysfunction (normal or mild organ dysfunction – TABLE 1C). DAF was scored as

"0" if the patient had organ dysfunction (moderate, severe, or extreme – TABLE 1C) or was not

alive during that 24-hour period. Each of the 28 days after ICU admission was scored in each patient in this fashion. Thus, the lowest score possible for each variable was zero and the highest score possible was 28. A low score is indicative of more organ dysfunction as there would be fewer days alive and free of organ dysfunction. There were no significant differences for the PAI-1 12580 genotypes between groups (either GG/GT vs TT or GG, GT and TT separately as shown in TABLES 4A and 4B below) with respect to age, sex, APACHE II score and percent of patients admitted with surgical diagnosis. Similarly, there were no significant differences for the PAI-1 4G/5G genotypes between groups (either 5G/5G vs 5G/4G and 4G/4G or any 4G separately as shown in TABLES 5A and 5B below).

	Age	% Female	APACHE II Score	% Surgical Diagnosis
GG	58 ± 2	38	24 ± 1	27
GT	57 ± 1	35	23 ± 1	20
TT	57 ± 1	41	23 ± 1	28
p Value	NS	NS	NS	NS

Table 4A. Baseline statistics of 587 Caucasian patients with SIRS grouped by PAI-1 12580 genotype. No significant differences in age, sex, APACHE II score or surgical diagnosis on admit were observed between PAI-1 12580 genotype groups (GG, GT or TT).

	Age	% Female	APACHE II Score	% Surgical Diagnosis
GG/GT	58 ± 1	36	23 ± 1	22
TT	57 ± 1	41	23 ± 1	28
p Value	NS	NS	NS	NS

Table 4B. Baseline statistics of 587 Caucasians with SIRS grouped by PAI-1 12580 genotype (GG/GT vs TT). No significant differences in age, sex, APACHE II score or surgical diagnosis on admit were observed between groups.

	Age	% Female	APACHE II Score	% Surgical Diagnosis
5G/5G	59 ± 2	42	22 ± 1	29
5G/4G	57 ± 1	34	22 ± 1	26
4G/4G	61 ± 2	33	21 ± 1	37
p Value	NS	NS	NS	NS

Table 5A. Baseline statistics of 314 Caucasians with SIRS grouped by PAI-1 4G/5G (837) genotype (5G/5G, 5G/4G and 4G/4G). No significant differences in baseline statistics were observed.

	Age	% Female	APACHE II Score	% Surgical Diagnosis
--	-----	----------	-----------------	----------------------

Any 4G	58 ± 1	34	22 ± 1	29
5G/5G	59 ± 2	42	22 ± 1	29
p Value	NS	NS	NS	NS

Table 5B. Baseline statistics of 314 Caucasians with SIRS grouped by PAI-1 4G/5G (837) allele (any 4G vs 5G/5G). No significant differences were observed in baseline statistics.

Patients having haplotypes containing the 12580 G allele had marked organ dysfunction of
 5 respiratory ($p < 0.05$), renal ($p < 0.05$), hepatic ($p < 0.05$), and neurologic ($p < 0.05$) organ systems
 over the 28 day study period (Figure 4). Days alive and free of all organ dysfunction for 12580
 GG was 1.8, GT 4.6, and TT 8.2 ($p < 0.01$). Thus, the 12580 G allele was associated with
 markedly worse overall organ failure. In addition, the 12580 G allele was associated with more
 SIRS ($p < 0.05$), more sepsis ($p < 0.05$), and more septic shock ($p < 0.05$) (Figures 5 and 6). Thus,
 10 the 12580 G allele was associated with markedly adverse clinical phenotype expressed as
 decreased survival, worse organ dysfunction, and worse SIRS.

It was found that there was no statistically significant difference or trend in DAF when grouped
 by PAI-1 4G/5G (837) genotype (Figures 8 A, 8B, 10 A, 10 B, 12 A, and 12 B). However, when
 15 patients were grouped by PAI-1 12580 genotype distinct trends in DAF were associated with
 increased numbers of the 12580G allele.

In Figure 7A, days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1
 12580 genotype (GG, GT and TT separately) in 587 critically ill Caucasians with the systemic
 20 inflammatory response syndrome (SIRS) is shown. Those patients possessing the 12580G allele
 had a strong trend toward fewer days alive and free of respiratory dysfunction and mechanical
 ventilation based on the number of copies of the 12580G allele they had. Those patients with 2
 copies of the 12580G allele ($N = 98$) had the fewest days alive and free of respiratory dysfunction
 (13 days) and mechanical ventilation (12 days). Patients with only one copy of the 12580G allele
 25 ($N = 284$) had intermediate values for days alive and free of respiratory dysfunction (13 days) and
 mechanical ventilation (13 days). Patients with 0 copies of the 12580 G allele (i.e. TT
 homozygotes; $N = 205$) had the greatest number of days alive and free of respiratory dysfunction
 (15 days) and mechanical ventilation (14 days).

Similarly, Figure 7B, shows days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 12580 genotype (GG/GT vs TT) in 587 critically ill Caucasians with SIRS. Those patients with either 1 or 2 copies of the 12580G allele (GG or GT, N = 382) had a strong trend toward fewer days alive and free of respiratory dysfunction ($p=0.057$) and mechanical ventilation ($p=0.077$) compared to those patients having 0 copies of the G allele (i.e. 12580 TT homozygotes, N = 205).

Figures 8A and 8B show days alive and free of respiratory dysfunction and mechanical ventilation by PAI-1 4G/5G (837) genotype (showing 4G/4G, 4G/5G and 5G/5G separately and any 4G vs 5G/5G) in 314 critically ill Caucasians with the systemic inflammatory response syndrome (SIRS). No significant differences or trends in DAF respiratory dysfunction or mechanical ventilation were observed based on PAI-1 4G/5G genotype.

In Figure 9A, days alive and free of acute lung injury (ALI) and P_aO_2/F_iO_2 (P/F) ratio greater than 300 mmHg by PAI-1 12580 genotype (GG, GT and TT separately) in 587 critically ill Caucasians with SIRS are shown. Those patients having the 12580G allele had a trend toward fewer days alive and free of ALI and $P/F < 300$ based on the number of copies of the 12580G allele they had. Patients with 2 copies of the 12580G allele (N = 98) had the fewest days alive and free of ALI (16 days) and $P/F < 300$ (5 days). Patients with 1 copy of the 12580G allele (N = 284) had intermediate values for days alive and free of ALI (16 days) and $P/F < 300$ (5 days). Patients with 0 copies of the 12580G allele had the greatest number of days alive and free of ALI (17 days) and $P/F < 300$ (7 days).

Similarly, Figure 9B, shows days alive and free of acute lung injury and $P_aO_2/F_iO_2 < 300$ mmHg by PAI-1 12580 genotype (GG/GT vs TT) for 587 Caucasians with SIRS. Those patients with either 1 or 2 copies of the 12580G allele (GG or GT, N = 382) had a trend toward fewer days alive and free of acute lung injury (16 days) and $PaO_2/FiO_2 < 300$ mmHg (5 days) compared to those patients with 0 copies of the 12580G allele (17 and 7 days, resp; N = 205).

Figure 10A shows days alive and free of acute lung injury (ALI) and P_aO_2/F_iO_2 (P/F) ratio greater than 300 mmHg by PAI-1 4G/5G/ (837) genotype (showing 4G/4G, 4G/5G and 5G/5G separately) in 314 critically ill Caucasians with SIRS. No significant differences or trends in DAF ALI or $P/F < 300$ were observed based on PAI-1 4G/5G genotype. Similarly, Figure 10B shows

days alive and free of acute lung injury and $P_aO_2/F_iO_2 < 300$ mmHg by PAI-1 4G/5G allele (any 4G vs 5G/5G) in 314 critically ill Caucasians with SIRS. No significant difference or trend in DAF ALI was observed based upon PAI-1 4G/5G allele. A trend toward fewer DAF P/F < 300 in the group "any 4G" was observed, but this trend was not as significant as that observed based on the PAI-1 12580 allele (Figure 9B).

In Figure 11A, days alive and free of steroid use by PAI-1 12580 genotype in 587 critically ill Caucasians with SIRS is shown. Patients with 2 copies of the 12580G allele ($N = 98$) had the fewest days alive and free of steroid use (15 days). Patients with 1 copy of the 12580G allele ($N = 284$) had similar values for days alive and free of steroid use (15 days). Patients with 0 copies of the 12580G allele ($N = 205$) had the greatest number of days alive and free of steroid use (17 days). Similarly, Figure 11B, shows days alive and free of steroid use by PAI-1 12580 genotype (GG/GT vs TT) for 587 critically ill Caucasians with SIRS. Those patients having either 1 or 2 copies of the 12580G allele (i.e. GG or GT; $N = 382$) had a strong trend toward fewer days alive and free of steroid use (15 days) compared to those patients with 0 copies (17 days; $N = 205$).

Figure 12A show days alive and free of steroid use by PAI-1 4G/5G (837) genotype in critically ill Caucasians with SIRS (grouped 4G/4G, 4G/5G and 5G/5G separately). No significant difference or trend in DAF steroid use was observed based on the PAI-1 4G/5G genotype.

Similarly, Figure 12B shows days alive and free of steroid use by PAI-1 4G/5G (837) were the same patients in 12A are grouped as any 4G vs 5G/5G. Again, no significant difference or trend was observed in DAF steroid use based on PAI-1 4G/5G allele.

Example 5 - Patient Outcomes for PAI-1 12580 Genotypes v. 4G/5G Genotypes in Surgery Patients

Plasminogen activator inhibitor-1 (PAI-1) levels were measured in 67 Caucasian cardiovascular surgery patients who had cardiopulmonary bypass. Serum levels were measured by enzyme-linked immunosorbent assay (ELISA) (Immulyse PAI-1, BioPool International) and presented based on PAI-1 12580 genotype (Figures 13A/B, 15A/B, 16 and 17A/B). No significant differences were observed between PAI-1 12580 genotype groups based on age or sex (as shown in TABLES 6A and 6B below).

	Age	% Female
GG	74 ± 5	60
GT	62 ± 2	27
TT	68 ± 2	33
p Value	NS	NS

Table 6A. Baseline statistics for 67 Caucasians who underwent on-pump cardiopulmonary bypass surgery and had PAI-1 serum levels measured. No significant differences in age or sex were observed between PAI-1 12580 genotype groups (GG, GT and TT).

	Age	% Female
GG/GT	64 ± 2	31
TT	68 ± 2	33
P Value	NS	NS

5 **Table 6B.** Baseline statistics for 67 Caucasians who underwent on-pump cardiopulmonary bypass surgery and had PAI-1 serum levels measured. No significant differences in age or sex were observed between PAI-1 12580 genotype groups (GG/GT vs TT).

10 Similarly, PAI-1 levels were also measured in 55 Caucasian cardiovascular surgery patients who had cardiopulmonary bypass and for comparison purposes serum levels were presented based on PAI-1 4G/5G (837) genotype (Figures 14A/B and 18A/B). A significant difference in proportion of female patients was observed (shown in TABLES 7A and 7B below), and this was accounted for in later analyses.

	Age	% Female
4G/4G	68 ± 2	53
4G/5G	64 ± 2	15
5G/5G	68 ± 2	36
p Value	NS	0.037

15 **Table 7A.** Baseline statistics for 55 Caucasians who underwent on-pump cardiopulmonary bypass surgery and had PAI-1 serum levels measured. No significant difference in age was observed between PAI-1 4G/5G groups (4G/4G, 4G/5G and 5G/5G). A significant difference in proportion of female patients was observed, and this was accounted for in later analyses.

	Age	% Female
Any 4G	66 ± 2	29
5G/5G	68 ± 2	36
p Value	NS	NS

Table 7B. Baseline statistics for 55 Caucasians who underwent on-pump cardiopulmonary bypass surgery and had PAI-1 serum levels measured. No significant differences in age or sex were observed between PAI-1 4G/5G genotype groups (any 4G vs 5G/5G).

5 Surprisingly, it was found that there was no statistically significant difference (or even a trend) in PAI-1 serum levels and systemic vascular resistance index (SVRI) in patients 4 hours post-operatively after cardiopulmonary bypass surgery when grouped by PAI-1 4G/5G (837) genotype. However, when patients were grouped by PAI-1 12580 genotype a statistically significant decrease in PAI-1 serum levels was associated with increased numbers of the 12580G allele, both
10 post and pre operatively. Similarly, the PAI-1 12580 genotype showed a statistically significant decrease in SVRI associated with the 12580G allele post-operatively. Thus, the PAI-1 12580 G allele was associated with more vasodilation (as shown by lower SVRI), and lower PAI-1 levels after cardiac surgery than PAI-1 12580 T. This represents independent confirmation in an independent cohort of patients who had cardiac surgery of the findings of PAI-1 12580 G in the
15 ICU cohort in that PAI-1 12580 G was associated with worse survival and organ dysfunction in the ICU cohort and lower SVRI and lower PAI-1 levels in the cardiac surgery cohort.

In Figure 13A, mean pre-operative PAI-1 serum levels in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG, GT and TT separately)
20 are shown. The 12580 TT (N = 3) homozygotes had a mean pre-operative PAI-1 serum level of 146.9 ng/mL, the GT heterozygotes (N = 14) had a mean pre-operative PAI-1 serum level of 102.2 ng/mL, and the GG homozygotes (N = 7) had a mean pre-operative serum PAI-1 level of 84.5 ng/mL. A statistically significant trend toward lower serum PAI-1 levels with increasing copy number of the 12580G allele is present ($p=0.044$). Statistics were performed on log
25 transformed data since levels were not normally distributed.

In Figure 13B, mean pre-operative serum PAI-1 levels in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG/GT vs TT) are shown. Here as before the TT homozygotes (N = 3) had higher mean pre-operative serum PAI-1 levels of
30 154.4 ng/mL compared to those patients with either the GG or GT genotypes (N = 21, mean serum PAI-1 level of 94.5 ng/mL). Statistical significance is not reached ($p=0.077$), but a value indicating a trend is observed. Statistics were performed on log transformed data.

In Figure 14A, serum PAI-1 levels measured 4 hours post-operatively in 55 Caucasians patients who underwent cardiopulmonary bypass surgery are shown. No significant difference was observed in serum PAI-1 level based on 4G/5G (837) genotype, and neither was there any trend in PAI-1 levels based on these genotypes. The difference in sex was included in a linear regression model and had no effect on lack of statistical significance. Similarly, Figure 14B shows serum PAI-1 levels measured 4 hours post-operatively in 55 Caucasians who underwent on-pump cardiopulmonary bypass surgery by any 4G vs. 5G/5G. Again, no significant difference in serum PAI-1 levels was observed based on 4G/5G (837) allele (any 4G vs 5G/5G), and neither was any significant trend observed. Statistics were performed on log-transformed data.

In Figures 15A and 15B, mean serum PAI-1 levels of 67 Caucasians who underwent on-pump cardiopulmonary bypass surgery collected at 4 hours post surgery grouped by PAI-1 12580 genotype (TT, GT and GG separately and GG/GT vs TT) are shown. This group includes as a subset those patients (N = 24) for whom data was collected at the pre-operative time-point, and therefore represents a confirmation of the trends observed in the aforementioned group (i.e. 12580G allele associated with lower serum PAI-1 levels) in a group with a larger sample size. The 12580 TT homozygotes (N = 21) had the highest mean serum PAI-1 levels at 195.1 ng/mL, followed by the GT heterozygotes with 130.3 ng/mL and the GG homozygotes with 111.3 ng/mL. In Figure 15B, the 12580 TT homozygotes had statistically significantly higher mean serum PAI-1 levels post-operatively (195.1 ng/mL) compared to those patients with either the GT or GG genotypes (126.0; $p=0.042$). A clear trend once again emerges wherein lower serum PAI-1 levels are associated with increasing numbers of the 12580G allele. Statistics were performed on log transformed data.

In Figure 16 mean differences in serum PAI-1 levels from pre-operative to 4 hours post-operative in 24 Caucasians who underwent on-pump cardiopulmonary bypass surgery is shown. Those patients who were homozygous for the 12580T allele had the greatest increase in PAI-1 levels pre-operative to post-operative (208.0 ng/mL). Heterozygous patients had an intermediate increase in PAI-1 levels (158.9 ng/mL), and those patients who were homozygous for the 12580G allele had the lowest mean increase in serum PAI-1 levels pre-operative to post-operative (43.6 ng/mL). Although statistical significance was not achieved, a clear trend is visible indicating a smaller increase in PAI-1 levels following cardiopulmonary bypass surgery with increasing copy number of the PAI-1 12580 G allele.

In Figure 17A, systemic vascular resistance index (SVRI) is shown 4 hours post-operatively in 66 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG, GT and TT separately). SVRI is calculated as $SVRI = (\text{mean arterial pressure} - \text{central venous pressure}) / \text{cardiac index}$, and expressed as $\text{dynes/s}\cdot\text{cm}^{-5}$. The PAI-1 12580 G allele is associated with a statistically significant trend toward lower mean SVRI 4 hours post-operatively based on the allele copy number: GG homozygotes have the lowest SVRI values ($1493.9 \pm 144.8 \text{ dynes/s}\cdot\text{cm}^{-5}$), GT heterozygotes have intermediate SVRI values ($1629.7 \pm 61.5 \text{ dynes/s}\cdot\text{cm}^{-5}$), and TT homozygotes have the highest SVRI values ($1780.4 \pm 64.6 \text{ dynes/s}\cdot\text{cm}^{-5}$). This trend is significant at the α level 0.05 ($p=0.028$).

Similarly, in Figure 17B, systemic vascular resistance index (SVRI) is shown 4 hours post-operatively in 66 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 12580 genotype (GG/GT vs TT). SVRI was calculated as described above for Figure 17A. The PAI-1 12580 G allele (here as GG homozygotes or GT heterozygotes) is shown to be significantly associated with lower mean SVRI 4 hours post-operatively than the T allele ($1604.4 \pm 57.8 \text{ dynes/s}\cdot\text{cm}^{-5}$ vs $1780.4 \pm 64.6 \text{ dynes/s}\cdot\text{cm}^{-5}$ respectively; $p = 0.050$).

Figure 18A shows the systemic vascular resistance index (SVRI) 4 hours post-operatively in 364 Caucasians undergoing on-pump cardiopulmonary bypass surgery grouped by PAI-1 4G/5G (837) genotype (4G/4G, 4G/5G and 5G/5G separately). SVRI is calculated as $SVRI = (\text{mean arterial pressure} - \text{central venous pressure}) / \text{cardiac index}$, and expressed as $\text{dynes/s}\cdot\text{cm}^{-5}$. No significant difference or trend in 4 hour post-operative SVRI was observed based upon the PAI-1 4G/5G genotype. Similarly, in Figure 18B in which patients were grouped by PAI-1 4G/5G (837) allele (any 4G vs 5G/5G). SVRI is calculated as above for 18A and again no significant difference or trend in 4 hour post-operative SVRI was observed based on the PAI-1 4G/5G allele.

Statistical Analysis

We compared measures of disease severity using dominant and co-dominant models. We tested for differences between genotype groups using ANOVA for continuous data and a Fisher exact test for discrete data. When appropriate Student's t-test and linear regression were also used.

The sample size described herein (260), compares with sample sizes in trauma of 61 (MENGES T et al. *Lancet* (2001) 357(9262):1096-7), in pediatric meningococcemia of 175 (DAWSON et al. (1993)), and adult meningococcal septic shock of 50 (WESTENDORP RG et al. *Lancet* (1999) 354(9178):561-3). Another possible source of error in association studies is population admixture (CARDON LR and BELL JI *Nature Reviews Genetics* (2001) 2:91-9). This issue was addressed by limiting the analysis to Caucasians and also because this group comprised the majority of our patients available for study.

SUMMARY

10 The PAI-1 12580 G allele is associated with greater evidence of SIRS and severe cardiovascular and respiratory dysfunction in a critically ill SIRS population and in a post CPB SIRS population. The critically ill SIRS population demonstrates that severe SIRS in the patients with the GG genotype was associated with more severe SIRS and more cardiovascular and respiratory failure (including more acute lung injury, more use of vasopressors, more use of steroids), but also in
15 trends to additional organ system dysfunction and importantly, to decreased survival. These observations were confirmed in an analogous but completely independent SIRS population of critically ill patients who had CPB. In the CPB population SIRS was induced by cardiac surgery and the cardiopulmonary bypass procedure itself without evidence of infection. Evidence for increased SIRS in those patients having the GG genotype in this population is provided by the
20 observation of greater reduction in systemic vascular resistance index (SVRI).

Critically ill patients with the 12580 G allele had significantly worse outcomes as indicated by lower survival, more SIRS, more severe cardiovascular and respiratory failure, and strong trends to more severe hepatic renal, neurologic, and coagulation dysfunction. The poor clinical
25 phenotype of the patients who had the 12580 G allele was also associated with greater use of corticosteroids. It is suspected that the reason for increased use of corticosteroids (see Figures 11A and B), is that the clinicians judged that there was a greater need for steroid treatment for severe shock and possibly prolonged respiratory failure.

30 Described herein are haplotypes that are novel in their ability to predict patient outcome due to inflammation, and haplotypes that are much more functional in describing patient risk than the previously described 4G/5G polymorphism. As a result, provided herein is a better strategy of predicting which patients are at greatest risk of an adverse outcome.

The PAI-1 4G allele is shown here to be divided into two distinct haplotype clades as a result of a remote recombination event. Inspection indicates that a recombinant event occurred between the 4G/5G site (position 837) in the promoter region and position 4588 (see Figure 1). In essence, a “natural experiment” occurred in which the promoter region containing the 4G allele in question, was recombined with a completely different haplotype clade (Figure 2). *Pan troglodytes* (chimpanzee) genotyping (Figure 2) suggests that 4G originated with the ancestral clade F (defined by the 12580 G allele) and recombined with the evolutionarily most distant haplotype clade D (defined by the 13985 T allele) of the PAI-1 gene (Figure 2). This “natural experimental design” therefore directly tests the hypothesis that the promoter region containing 4G is causal because the 4G allele is distributed into two different haplotype clades that can then be compared. It was found that the two 4G haplotype clades (F and D) were associated with very different outcomes. The 12580 G haplotype clade was associated with significantly reduced hospital survival and organ dysfunction while the 13985 haplotype clade was not. Thus, this “natural experiment” has lead to the conclusion that 4G is not associated with patient outcome or not strongly so.

From our results the 4G allele of PAI-1 is not associated with poor prognosis. As described herein a SNP within the 12580 G haplotype clade (clade F) is a better candidate for predicting patient outcome, because the 12580 G haplotype clade was strongly associated with adverse outcomes. In the haplotype map (Figure 1), the PAI-1 4G allele (837-) and 12580 G allele are linked by some measure of linkage disequilibrium, though not complete LD (estimated to be about $r^2 \approx 0.2$ calculated using the Visual Genotype2 program available at <http://droog.gs.washington.edu/pgs/VG2.html>). As a result the 12580G allele is proposed herein as a better predictor of patient outcome than the 4G allele.

In a cohort of critically ill patients, 95% having SIRS, analysis based on the haplotype structure of the PAI-1 gene suggested that the 4G/5G polymorphism was not associated with adverse outcome. Rather, a polymorphism contained within the haplotype clade defined by the minor G allele of the 12580 T/G polymorphism, was a better candidate. The 12580 G haplotype clade was associated with decreased survival and increased organ dysfunction, severity of SIRS, and occurrence of sepsis and septic shock. The association of the 12580 G haplotype clade with adverse outcome is important because it shows that the very commonly reported 4G/5G SNP is

not the best predictor of patient outcome. This has implications for critically ill patients who have SIRS and possibly for patients having other diseases (e.g. stroke, myocardial infarction) in which 4G/5G has been reported as associated with adverse outcomes.

- 5 The PAI-1 12580 G allele was associated with lower serum PAI-1 levels (Figures 13A/B and 15A/B) and with fewer days alive and free of organ dysfunction (respiratory, acute lung injury), support (mechanical ventilation and steroid use) and is associated with significantly lower systemic vascular resistance index in cardiac surgical patients. Therefore, the 12580 G allele is a novel indicator of poor outcome (survival and days alive and free of organ dysfunction as
- 10 described above) and lower serum PAI-1 levels.

In conclusion, the 4G/5G polymorphism of PAI-1 is not as good a predictor of outcome in patients with SIRS, but rather, a polymorphism contained within the haplotype clade defined by the minor G allele of the 12580 T/G polymorphism, is a better candidate.

CLAIMS

What is Claimed is:

- 5 1. A method for obtaining a prognosis for a patient having or at risk of developing an inflammatory condition, the method comprising determining a genotype for one or more polymorphism sites in the plasminogen-activator-inhibitor-1 (PAI-1) gene for the patient, wherein said genotype is indicative of an ability of the patient to recover from an inflammatory condition, provided that the one or more polymorphism sites is not solely a
10 polymorphism at a site corresponding to position 837 of SEQ ID NO.: 1.
2. The method of claim 1, wherein the one or more polymorphism sites includes position 12580 of SEQ ID NO.: 1 or a polymorphism site in total linkage disequilibrium thereto.
- 15 3. The method of claim 2, wherein the one or more polymorphism sites are selected from positions 5645, 7121, 7437, 8070, 8406, 9463, 9466, 12219, 12580, 13889 and 14440 in SEQ ID NO: 1.
4. The method of claim 1, wherein the genotype comprises a combination of two or more
20 polymorphism sites, the combination being selected from the group of positions corresponding to SEQ ID NO:1 consisting of:
 - 664 A and 2037 T;
 - 664 A and 2362 -;
 - 664 A and 2852 A;
 - 25 664 A and 5834 A;
 - 837 - and 2037 T;
 - 837 - and 2362 -;
 - 837 - and 2852 A;
 - 837 - and 5834 A;
- 30 one of 5878 G and one of 7365 T and one of 4588 T
 7343 G 7729 + 5404 G
 13605 A 7771 A 5686 A
 12750 A 5984 A
 53

11312 A; and

one of 2846 A and 10381 T and one of 7365 T and one of 4588 T
 6821 T and 10381 T 7729 + 5404 G
 5 9759 G and 10381 T 7771 A 5686 A
 12750 A 5984 A
 11312 A.

5. The method of any one of claims 1-4, further comprising comparing the genotype so
 10 determined with known genotypes, which are indicative of a prognosis for recovery from
 the same inflammatory condition as for the patient or another inflammatory condition.
6. The method of any one of claims 1-5, further comprising obtaining a plasminogen-
 activator-inhibitor-1 gene sequence of the patient.
7. The method any one of claims 1-5, wherein said determining of genotype is performed on
 a nucleic acid sample from the patient.
8. The method of claim 7, further comprising obtaining a nucleic acid sample from the
 20 patient.
9. The method any one of claims 1-8, wherein said determining of genotype comprises one
 or more of:
 - (a) restriction fragment length analysis;
 - 25 (b) sequencing;
 - (c) hybridization;
 - (d) oligonucleotide ligation assay;
 - (e) ligation rolling circle amplification;
 - (f) 5' nuclease assay;
 - 30 (g) polymerase proofreading methods;
 - (h) allele specific PCR; and
 - (i) reading sequence data.

10. The method of any one of claims 1-9, wherein the genotype of the patient is indicative of a decreased likelihood of recovery from an inflammatory condition.
11. The method of claim 10, wherein the prognosis is indicative of severe cardiovascular or respiratory dysfunction in critically ill patients.
12. The method of claim 10 or 11, wherein the genotype is selected from the group of single polymorphism sites and combined polymorphism sites consisting of:
- 5645 T;
 - 7121 G;
 - 7437 T;
 - 8070 A;
 - 8406 C;
 - 9463 G;
 - 9466 T;
 - 12219 C;
 - 12580 G;
 - 13889 C;
 - 14440 A;
 - 664 A and 2037 T;
 - 664 A and 2362 -;
 - 664 A and 2852 A;
 - 664 A and 5834 A;
 - 837 - and 2037 T;
 - 837 - and 2362 -;
 - 837 - and 2852 A; and
 - 837 - and 5834 A.
13. The method of any one of claims 1-9, wherein the genotype of the patient is indicative of an increased likelihood of recovery from an inflammatory condition.
14. The method of claim 13, wherein the prognosis is indicative of less severe cardiovascular or respiratory dysfunction in critically ill patients.

15. The method of claim 13 or 14, wherein the genotype is selected from the group of single polymorphism sites and combined polymorphism sites consisting of:

5645 C;

7121 A;

7437 C;

8070 G;

8406 T;

9463 A;

9466 C;

12219 T;

12580 T;

13889 T;

14440 G;

15 one of 5878 G and one of 7365 T and one of 4588 T
 7343 G 7729 + 5404 G
 13605 A 7771 A 5686 A
 12750 A 5984 A
 11312 A; and

20 one of 2846 A and 10381 T and one of 7365 T and one of 4588 T
 6821 T 7729 + 5404 G
 9759 G 7771 A 5686 A
 12750 A 5984 A
 11312 A.

16. The method of any one of claims 1-15, wherein the inflammatory condition is selected from the group consisting of: sepsis, septicemia, pneumonia, septic shock, systemic inflammatory response syndrome (SIRS), Acute Respiratory Distress Syndrome (ARDS), acute lung injury, infection, pancreatitis, bacteremia, peritonitis, abdominal abscess, inflammation due to trauma, inflammation due to surgery, chronic inflammatory disease, ischemia, ischemia-reperfusion injury of an organ or tissue, tissue damage due to disease,

tissue damage due to chemotherapy or radiotherapy, and reactions to ingested, inhaled, infused, injected, or delivered substances, glomerulonephritis, bowel infection, opportunistic infections, and for patients undergoing major surgery or dialysis, patients who are immunocompromised, patients on immunosuppressive agents, patients with HIV/AIDS, patients with suspected endocarditis, patients with fever, patients with fever of unknown origin, patients with cystic fibrosis, patients with diabetes mellitus, patients with chronic renal failure, patients with bronchiectasis, patients with chronic obstructive lung disease, chronic bronchitis, emphysema, or asthma, patients with febrile neutropenia, patients with meningitis, patients with septic arthritis, patients with urinary tract infection, patients with necrotizing fasciitis, patients with other suspected Group A streptococcus infection, patients who have had a splenectomy, patients with recurrent or suspected enterococcus infection, other medical and surgical conditions associated with increased risk of infection, Gram positive sepsis, Gram negative sepsis, culture negative sepsis, fungal sepsis, meningococcemia, post-pump syndrome, cardiac stun syndrome, myocardial infarction, stroke, congestive heart failure, hepatitis, epiglottitis, E. coli 0157:H7, malaria, gas gangrene, toxic shock syndrome, mycobacterial tuberculosis, Pneumocystis carinii, pneumonia, Leishmaniasis, hemolytic uremic syndrome/thrombotic thrombocytopenic purpura, Dengue hemorrhagic fever, pelvic inflammatory disease, Legionella, Lyme disease, Influenza A, Epstein-Barr virus, encephalitis, inflammatory diseases and autoimmunity including Rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, inflammatory bowel disease, idiopathic pulmonary fibrosis, sarcoidosis, hypersensitivity pneumonitis, systemic vasculitis, Wegener's granulomatosis, transplants including heart, liver, lung kidney bone marrow, graft-versus-host disease, transplant rejection, sickle cell anemia, nephrotic syndrome, toxicity of agents such as OKT3, cytokine therapy, and cirrhosis.

17. The method of any one of claims 1-16, wherein the inflammatory condition is systemic inflammatory response syndrome.

18. A method of identifying a polymorphism in a PAI-1 gene sequence that correlates with a patient prognosis, the method comprising:

- a) obtaining PAI-1 gene sequence information from a group of patients;
- b) identifying a site of at least one polymorphism in the PAI-1 gene;

- c) determining genotypes at the site for individual patients in the group;
- d) determining an ability of individual patients in the group to recover from the inflammatory condition; and
- e) correlating genotypes determined at step (c) with patient abilities determined at step (d).

19. The method of claim 18, wherein the inflammatory condition is selected from the group consisting of: sepsis, septicemia, pneumonia, septic shock, systemic inflammatory response syndrome (SIRS), Acute Respiratory Distress Syndrome (ARDS), acute lung injury, infection, pancreatitis, bacteremia, peritonitis, abdominal abscess, inflammation due to trauma, inflammation due to surgery, chronic inflammatory disease, ischemia, ischemia-reperfusion injury of an organ or tissue, tissue damage due to disease, tissue damage due to chemotherapy or radiotherapy, and reactions to ingested, inhaled, infused, injected, or delivered substances, glomerulonephritis, bowel infection, opportunistic infections, and for patients undergoing major surgery or dialysis, patients who are immunocompromised, patients on immunosuppressive agents, patients with HIV/AIDS, patients with suspected endocarditis, patients with fever, patients with fever of unknown origin, patients with cystic fibrosis, patients with diabetes mellitus, patients with chronic renal failure, patients with bronchiectasis, patients with chronic obstructive lung disease, chronic bronchitis, emphysema, or asthma, patients with febrile neutropenia, patients with meningitis, patients with septic arthritis, patients with urinary tract infection, patients with necrotizing fasciitis, patients with other suspected Group A streptococcus infection, patients who have had a splenectomy, patients with recurrent or suspected enterococcus infection, other medical and surgical conditions associated with increased risk of infection, Gram positive sepsis, Gram negative sepsis, culture negative sepsis, fungal sepsis, meningococcemia, post-pump syndrome, cardiac stun syndrome, myocardial infarction, stroke, congestive heart failure, hepatitis, epiglottitis, E. coli 0157:H7, malaria, gas gangrene, toxic shock syndrome, mycobacterial tuberculosis, Pneumocystic carinii, pneumonia, Leishmaniasis, hemolytic uremic syndrome/thrombotic thrombocytopenic purpura, Dengue hemorrhagic fever, pelvic inflammatory disease, Legionella, Lyme disease, Influenza A, Epstein-Barr virus, encephalitis, inflammatory diseases and autoimmunity including Rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, inflammatory bowel disease, idiopathic pulmonary fibrosis, sarcoidosis,

hypersensitivity pneumonitis, systemic vasculitis, Wegener's granulomatosis, transplants including heart, liver, lung kidney bone marrow, graft-versus-host disease, transplant rejection, sickle cell anemia, nephrotic syndrome, toxicity of agents such as OKT3, cytokine therapy, and cirrhosis.

5

20. A kit for determining a genotype at a defined nucleotide position within a polymorphism site in a PAI-1 gene sequence from a patient to provide a prognosis of the patient's ability to recover from an inflammatory condition, the kit comprising, in a package a restriction enzyme capable of distinguishing alternate nucleotides at the polymorphism site or a
10 labeled oligonucleotide having sufficient complementary nucleotides to an adjacent sequence at or near the polymorphism site and capable of distinguishing said alternate nucleotides, provided that the polymorphism site is not solely a polymorphism at a site corresponding to position 837 of SEQ ID NO.: 1.
- 15 21. The kit of claim 20, where the polymorphism site corresponds to one or more of positions 5645, 7121, 7437, 8070, 8406, 9463, 9466, 12219, 12580, 13889 and 14440 of SEQ ID NO: 1.
22. The kit of claim 21, where the polymorphism site corresponds to position 12580.
- 20 23. The kit of claim 20, 21 or 22 comprising said restriction enzyme and an oligonucleotide or a set of oligonucleotides suitable to amplify a region surrounding the polymorphism site.
24. The kit of claim 23, further comprising a polymerization agent.
- 25 25. The kit of any one of claims 20-24, further comprising instructions for using the kit to determine genotype.
- 30 26. A method for selecting a group of patients for determining the efficacy of a candidate drug known or suspected of being useful for the treatment of an inflammatory condition, the method comprising determining a genotype for one or more polymorphism sites in the plasminogen-activator-inhibitor-1 gene for each patient, wherein said genotype is indicative of the patient's ability to recover from the inflammatory condition and sorting

patients based on their genotype, provided that the polymorphism site is not solely a polymorphism at a site corresponding to position 837 of SEQ ID NO.: 1.

27. The method of claim 26 further comprising, comparing patient response to the candidate
5 drug based on genotype of the patient.
28. The method of claim 27, wherein patient response is determined by each patient's ability
to recover from the inflammatory condition.

1/18

Clade	Relative Position	Frequency
1	12860	2
2	12405	2
3	12309	1
4	12025	2
5	11737	3
6	11560	1
7	11473	1
8	11267	8
9	11170	5
10	11000	2
11	10639	2
12	9732	1
13	8801	1
14	8179	1
15	7886	1
16	7883	1
17	6931	1
18	6826	1
19	6512	1
20	6490	1
21	6191	1
22	6149	1
23	5857	1
24	5849	1
25	5785	1
26	5763	1
27	5541	1
28	5241	1
29	4404	1
30	4298	1
31	4254	1
32	4106	1
33	4065	1
34	3824	1
35	3557	1
36	3008	1
37	1928	1
38	1868	1
39	1828	1
40	1824	1
41	1688	1
42	1272	1
43	1266	1
44	782	1
45	457	1
46	-743	1
47	-916	1
48	-1168	1

FIGURE 1

2/18






FIGURE 2

3/18

FIGURE 3




4/18

FIGURE 4


5/18

FIGURE 5



6/18

FIGURE 6



7/18

FIGURE 7A





FIGURE 7B



8/18

FIGURE 8A





FIGURE 8B



9/18

FIGURE 9A





FIGURE 9B



10/18

FIGURE 10A





FIGURE 10B



11/18

FIGURE 11A





FIGURE 11B



12/18

FIGURE 12A





FIGURE 12B



13/18

FIGURE 13A





FIGURE 13B



14/18

FIGURE 14A





FIGURE 14B



15/18

FIGURE 15A






FIGURE 15B



16/18

FIGURE 16



17/18

FIGURE 17A





FIGURE 17B



18/18

FIGURE 18A





FIGURE 18B



SEQUENCE LISTING

<110> THE UNIVERSITY OF BRITISH COLUMBIA
 RUSSELL, James A.
 WALLEY, Keith R.

<120> PLASMINOGEN ACTIVATOR INHIBITOR-1 (PAI-1) HAPLOTYPES USEFUL AS
 INDICATORS OF PATIENT OUTCOME

<130> 80021-638

<140> NOT YET ASSIGNED

<141> 2004-03-19

<150> US 60/455,550

<151> 2003-03-19

<160> 11

<170> PatentIn version 3.2

<210> 1

<211> 14544

<212> DNA

<213> Homo sapiens

<400> 1
 cgtactggtc catagcctgt taggaaccag gctgcataac aggaggtgag tggcaggtga 60
 gtgaaatttc atctgtagtt acagccactc ctcatcactc gcattaccac cagagctcca 120
 ctccctgtca gatcagcggc ggcattagat tctcatagga gctcgaaccc tattctaaac 180
 tgttcatgtg agggatctag gttgcaagct ccctatgaga atctaagcc tgatgatctg 240
 tcacggtctc ccatcacccc tagatgggac catctagttg caggaaaaca agctcagggc 300
 tcccactgat tctacacgat ggtgaattgt ggaattatct cattatataat attacaatgt 360
 aataataata gaaataaagc acacaataaa tgtaattgtc ttgaatcatc ccgaaaccat 420
 cccaccctgg tctgtgaaaa aattgtcttc catgaaacca gtccctggtg ccaaaaacgt 480
 tgaggaccac tgtctcacag aatctatcgg tcaactcttc tccctcacc cccttgccct 540
 aaaagcacac cctgcaaacc tgccatgaat tgacactctg tttctatccc ttttcccctt 600
 gtgtctgtgt ctggaggaag aggataaagg acaagctgcc ccaagtccta gcgggcagct 660
 cgaggaaagt aaacttacac gttggtctcc tgtttcctta ccaagctttt accatggtaa 720
 cccctggtcc cggttcagcca ccaccacccc acccagcaca cctccaacct cagccagaca 780
 aggttgttga cacaagagag ccctcagggg cacagagaga gtctggacac gtgggggagt 840
 cagccgtgta tcatcggagg cggccgggca catggcaggg atgagggaaa gaccaagagt 900
 cctctgtttg gcccaagtcc tagacagaca aaacctagac aatcacgtgg ctggctgcat 960
 gccctgtggc tgttgggctg ggcccaggag gagggagggg cgctctttcc tggaggtggt 1020
 ccagagcacc ggggtggacag ccctggggga aaacttccac gttttgatgg aggttatctt 1080
 tgataactcc acagtgcctt ggttcgcca aaggaaaagca ggcaacgtga gctgtttttt 1140
 ttttctccaa gctgaacact aggggtccta ggcttttttg gtcacccggc atggcagaca 1200
 gtcaacctgg caggacatcc gggagagaca gacacaggca gagggcagaa aggtcaaggg 1260
 aggttctcag gccaaggcta ttggggtttg ctcaattgtt cctgaatgct cttacacacg 1320
 tacacacaca gagcagcaca cacacacaca cacacacatg cctcagcaag tccagagag 1380
 ggaggtgtcg agggggaccc gctggctgtt cagacggact ccagagcca gtgagtgggt 1440
 ggggctggaa catgagttca tctatttctt gccacatctt ggtataaaaag gaggcagtgg 1500
 cccacagagg agcacagctg tgtttggctg cagggccaag agcgtgttca agaagaccca 1560
 cacgcccccc tccagcagct gaattcctgc agctcagcag ccgcccagc agcaggacga 1620
 accgccaatc gcaaggcacc tctgagaact tcaggtagga gaaaagcaaa ctccctccaa 1680
 cctcttactt cgggcttaag gcagagaact cgctcccca gaatctcctc cctccatgat 1740
 cccccgctat tctctatatt tcttttcttc ggacctgcag ccttgggtcg accctgacct 1800
 aggggtgact gcaggagagc agggaggatg gtcaggcgtc accaacaacc ccatcaccca 1860
 gtaacaagaa ccttgactct ctcagtcctt ctgcatcaag acacttacc atttccacc 1920
 tcatgcctgc taaactgaat gaaacaatcg ctgggaaagc attaagagaa ttaaggctgg 1980
 gcactgtggc tcatgcctgt aatcccagca ctttgtgagg ctgaggcagg cagataactt 2040

gagcccagga	gtttgagacc	agcctgggca	acatggcaaa	accctgctct	cccaaaaaaa	2100
tacaaaaatt	agctgggctg	gctgggtgtg	ctgtattccc	agctacttgg	gaggctgagg	2160
tgggaggatt	gcttcagctg	gggaggcgga	ggctgcaggg	agccaagact	gagccattgc	2220
accagcctg	ggtgacagag	caagaccctg	tctctaaaaa	tgaatgaaag	gaaggaagaa	2280
agagagagaa	agagagagag	gaaagaagga	aggaagtaaa	gaagaaagaa	agaaagaaag	2340
aggaaagagg	aagaaagaaa	gaaaagaaa	aaaagaaaa	aaagcaaatt	taaagcttat	2400
gcaaatcaaa	gatgttgtga	taattgataa	ttgagtctgg	gctaaattcc	ccctgggctg	2460
caaaggcaga	gagtggtaat	gacttctcac	ctgcttttct	tctaaggctt	ttttacggga	2520
cacagagggg	agggagatgg	actggattcc	aagattccca	cagggaaga	tgggcgaaga	2580
ctccctgcc	ctgcccgggg	ataagtcagt	ctgagtgaga	cggagtggga	tgggcttaga	2640
acctgaacat	gtcatggtct	cttcctgcac	cttgccctag	tgttcaacta	ccacctgctt	2700
gcaggaaaca	agaagagcag	ggcccacagc	tggccagctc	ccctccctc	ccgcctgtct	2760
tccagaacga	ttccttcacc	agccctcttt	ccattgctct	aggatgcaga	tgtctccagc	2820
cctcacctgc	ctagtccctg	gcctggccct	tgtcttttgt	gaagggctct	ctgtgcacca	2880
tccccatcc	tacgtggccc	acctggcctc	agacttcggg	gtgagggtgt	ttcagcaggt	2940
ggcgaggcc	tccaaggacc	gcaacgtggt	tttctcacc	tatggggtgg	cctcgtgtgt	3000
ggccatgctc	cagctgacaa	caggaggaga	aaccagcag	cagattcaag	cagctatggg	3060
attcaagatt	gatggtgagc	cacgggacac	caggggaggt	gggtggcatg	cagaacagac	3120
ctaccagaag	ccaaggaaag	gctggctctg	gcttagccga	gccaagcccc	atacagctgt	3180
gctgcagggg	ccaccccatc	ttcttccac	tacactccaa	gtcactggac	ccttgaatct	3240
ccaaggggtg	ctgaccagta	gatttaccgc	ttattcacca	ccgtgtgatc	ttaacctcgt	3300
taagtttgcc	catctacaaa	atgaggatta	tttgctgtcc	taaagaattc	atgagccggg	3360
cgcggtggct	caaacgcctg	taatcccagc	actttgggag	gccaaggcgg	gcggtatcat	3420
aggtcaggag	atcaagacca	tcctggctaa	cacagtgaag	ctccatctct	actaaaaata	3480
caaaaaaat	tagccaggcg	tggtggcagg	cgctgtagt	ccagctact	cgggagctgt	3540
aggcaggaga	atggcatgaa	cccaggaggc	agagcttgca	gtgagctgag	atcgtgccac	3600
tgactccag	cctgggcgac	agagagagac	tccgtctcaa	aaaaaaaaat	taaaagaatt	3660
catggaatta	cacttgtgaa	atacttagca	tagccatcac	tataggaaaa	aatctaaagg	3720
ccaggcacag	tgcctcatgc	ctgtaatctc	agcactttcg	gagtttgagg	caggaggatc	3780
acccaaggct	aggagttaa	ggccagcctg	ggcaatacgg	tgaacccccg	tctctaataa	3840
aaatataaaa	attagtctga	tgagggtggg	caactgtaat	cccagctact	caggaagctg	3900
agacacaaga	atcactttta	cccgggaggt	ggagggtgga	gtgagctgag	atcacaccat	3960
tgcactccag	cctgggtgac	agagttagac	ctgtcaaaaa	aaagaaaaga	aagagagaga	4020
gagagagaag	agagagaaag	aaagaagaag	aaagaagaag	agagagagag	agaaagaaag	4080
aaagaaacaa	agaaacaaag	aaagaaagga	aaagaaaaaa	aaaactaagg	ccaggcaagg	4140
tggttatga	ctgtaatttc	agcacttttg	aagattgagg	caggaggatc	acttgaggcc	4200
agaagtgcga	gacaagactg	agcaacaggg	agaccctgc	ctctacaaaa	aaatttacaa	4260
attagccaga	tgtggtgaca	catacctgta	gtcccaacta	ctcaggaggc	tgagggtggga	4320
ggatggcttg	agcccaggag	ctggaggctg	cagttagcta	tgattgtacc	actgcacttc	4380
agcctgggca	acaaagggaa	gccctgtctg	aaaaaaaaaa	aaaaagaaaa	agaagaagaa	4440
agaaaatatt	tagggttcat	ccaggaggga	gagggtgcag	taagctgaca	tcgcgccatt	4500
gcactccagc	ctgggagaca	agagcaaaac	tccaactcaa	aaaaaaaaaa	aaaaaaaaaa	4560
caggaaagaa	atatttaggg	ttcataactt	aagaacagag	aaaaatattc	tagcccaaag	4620
aaaggggttg	gatctgagac	ttttgaagaa	aggaaggaga	tacagaaaag	agatttcac	4680
ctggaatgaa	atctccctcc	agagagccct	gggaaagcac	ggtagccccc	atccatcaga	4740
gtggagcccc	ttgtggggga	agtgggctcg	gctgggaacc	ctcaattcag	cataagcctc	4800
acatgtctc	tcctctctgt	cccgggtgcg	acaagggcat	ggcccccgcc	ctccggcatc	4860
tgtacaagga	gctcatgggg	ccatggaaca	aggatgagat	cagcaccaca	gacgcgatct	4920
tcgtccagcg	ggatctgaag	ctggtccagg	gcttcatgcc	ccacttcttc	aggctgttcc	4980
ggagcacggg	caagcaagtg	gacttttcag	aggtggagag	agccagattc	atcatcaatg	5040
actgggtgaa	gacacacaca	aaaggtgagc	agcaggggaa	aggaaccca	tttctgtggc	5100
ctcaagagaa	agggaatttg	gaaataaatc	cacatatccc	agttgggtgc	agtagttcac	5160
acctgtaatc	ccagcccaac	actttgggag	gtctaggcga	gaggaaggct	tgaggcctgg	5220
agtttgagac	cagcctggcc	aacataacaa	gacctcatct	cttcaaaaaa	tttaaaaacc	5280
agccgggcat	ggtggtgcac	acctgtagtc	ccagctactt	gggaggctga	ggtgggagga	5340
tcacttgagt	ccagcagttc	aaggctgcag	tgagctatgt	ttgcaccacc	acactccagc	5400
ctgagtcaca	gaacaagacc	tcatctctaa	aaaacaaaca	aaaaccaa	ccacatatcc	5460
taaaaaatgc	tccttttcag	cattctcttc	tctatggaca	aagggctgga	tgctttaaga	5520
accaaattct	aggctgggca	cggtggctca	cgctctaat	cctagcactt	tgagaggcca	5580
aggcggtgag	attgcctgag	cacaggaggt	cgagaccagc	ctggccaaca	tggtgaaacc	5640
ctgtctctgt	caaaaataca	aaaaattagc	cagggtgtgt	ggcgctgcc	tataatccca	5700

gctgctcggg	aggatgaggt	tcaaagaatc	acttgaaccc	gggaggcaga	ggctgcagt	5760
agctgagatc	atgccactgc	actccagcct	gggtgacaga	gcaagacttt	gtctccaaaa	5820
aaaggaacta	gacgggttca	tttaaaccce	tgactgcagc	cctttgacat	acatccaatt	5880
gaggactggg	gactccggga	aacatctaaa	aggcttaaaa	actttgtcta	acttcagecg	5940
ggcatggtgg	ctcacacctg	taatcccagc	actttgggag	gctgaggcag	gtggatcaca	6000
aggtcaggag	tttgagacga	gcctgaccaa	catggtgaaa	ccccgtctct	actaaaaata	6060
caaaaattag	ccaggcatgg	tggcaggcgc	ctgtaatccc	agctattcgg	gaggctgagg	6120
caggagaatt	gcttgaaccc	cggagacaga	ggttgacagc	agccgagatc	gcgccactgc	6180
actccagcct	ggcaatagag	tgagactcca	tctcaaaaaca	acaacaacaa	caacaacaac	6240
aacaaaatcg	tctaacttcc	tgatcttcc	gatcattgat	tttcccatag	gtatgatcag	6300
caacttgctt	gggaaaggag	ccgtggacca	gctgacacgg	ctggtgctgg	tgaatgccct	6360
ctacttcaac	ggccagtggg	agactccctt	ccccgactcc	agcaccacc	gccgcctctt	6420
ccacaaatca	gacggcagca	ctgtctctgt	gccccatgat	gctcagacca	acaagttcaa	6480
ctatagtaag	tccaagagcc	ccttccccac	agcccacagc	aactgcatct	cattcctggg	6540
gtctcccaag	gaatacccaa	aatgtcaccc	tctgaggagg	gaagaccaca	gggaatgctc	6600
ccctttaagg	gaggagagac	cctagaatat	actccagctt	tgacaaagat	ttcccaagca	6660
ggagacatca	ggataatggg	aacagaagac	aggagggtta	tcccatgaag	gatgaagaag	6720
ctgaaatcca	gagattccct	cagggccaca	tttgtccacc	tgactccagg	gtctcatctt	6780
cgtgtgttgc	tagtgtgatt	acctggggat	gagaaatcct	gctgggggag	ttgagggttaa	6840
gaggatgagg	actccagggtg	ctgtggctca	cgctgtaat	cccagcactt	tgggaggcca	6900
aggcagggtg	atcaggagtt	tgaggtcagg	agtttgagac	cagcctggcc	aacatggtga	6960
aaccctgtct	ctactaaaaa	tgcaaaaatt	agccagggtg	ggtggcaggc	gcctgtaatc	7020
ccagctactc	gggaggctga	ggcaggagaa	tcacttgagc	ccgggagggtg	gaggttgcat	7080
tgagccgaac	gaaattgagc	cacttcaccc	cagcctgggc	aaaagagtga	aattccattc	7140
aaaaaataaa	aaaaaataaa	aaggatgagg	actgggtggc	actgggtggc	gggtgtgggg	7200
aaaatggaag	tgaaggaagg	ccaaaagaga	cagagaaggc	ctggcgcggc	gactcacgcc	7260
tataatccca	gcactttggg	aggctgagaa	gggggattgc	ttgaggccag	aagttgaata	7320
ccagtctggg	cagcatagca	agaccctgcc	tctacaaaaa	aaaaattttt	tttaattagc	7380
caggcttggt	gacatgcac	tgtagtctac	tcaagaagct	gaggtgaggc	caggcacggt	7440
ggctcacgcc	tgtattccca	gcactttggg	aggccaaggc	gggtggatga	cctgaggtca	7500
ggagttcaag	accagcctgg	ccaacatggt	gaaaccccat	ctgtataaaa	atacaaaaat	7560
tagctgggca	tgatagcagg	tgccgtgaat	tccagctact	caggaggctg	aggtgggaga	7620
atctattgaa	cccgggaggg	ggagggtgca	gtgagccgag	atcatgccat	tgcaactccag	7680
cctgggcgac	agagtgcagc	tccttctcaa	aacaaacaaa	caaacaacaa	aacaaaatac	7740
agaagctgag	gcgggaggaa	catttgaaac	ggattcggag	gctgcagtga	gctatgattg	7800
caccactgcg	ctccagtctg	tgtgacagtg	agaccctgtc	tcttacacac	acacacacac	7860
acacacacac	acatgcacac	acacagagag	agagaaatta	gaagatactg	aattggcaga	7920
agagaaggga	aatagaatt	aaaatactga	ataggggagc	agtgaacagg	ggatacccaa	7980
aagccaagag	cgagagagag	cctggcttcc	agaaatagtg	gagaagccag	gagaactagg	8040
tgaaaaccca	gtgctgggtt	gccatcagcg	agagctggag	ccatttccaa	cgaaccatct	8100
tgtcgtcttc	acagctgagt	tcaccacgcc	cgatggccat	tactacgaca	tcctggaact	8160
gccctaccac	ggggacaccc	tcagcatggt	catgtctgcc	ccttatgaaa	aagaggtgcc	8220
tctctctgcc	tcaccaacaa	ttctgagtgc	ccagctcatc	agccactgga	aaggcaacat	8280
gaccaggctg	ccccgcctcc	tggttctgcc	caagtaagcc	accccgtat	ctccccgacc	8340
taccaacccc	tctctcctgg	ctccctaaag	tcaccgcccc	caggttgaat	ttcccagatc	8400
tgtgatgctt	gcaggacatg	catgtgtggg	aggctgatgg	gaaactgtgg	cctgggtttg	8460
attatgagtc	ttgcaatcat	ccctccccct	gttctctgctg	gagggcaggg	gacagctctt	8520
cctgaccaca	ccccacatt	gactatcccc	agaataccca	gcaaaagccc	ccaaaaggag	8580
agtcagagaa	atgaggagg	tgggggcccc	atcagtcac	atctacttag	ggtcgcccca	8640
tcagcacttc	catccccaac	cctttcaagt	caacatccaa	acaaaagaaa	tcacttccaa	8700
ggagggagca	gtcacaagcg	cagcttctag	ctggggttcc	aagaaagcag	atttttcgaa	8760
atccttctgc	agaaggaagc	aaagagattt	tttgaaatct	ttctgcagaa	ggagaaggct	8820
ggagctgggg	aactccagaa	ttatagggaa	gcctcccacc	acgctcatcc	caaatttccg	8880
gatgctataa	tgccaggctt	ggggaaagag	gagaatttag	ttggtttagct	ggtgcgtgct	8940
ctcacttgca	tcctctctct	tcctcttttt	ttttttctct	ctctctctct	ggctcataaa	9000
aatggaggta	attagttgtg	ccctggtgag	aagcagagag	tgacaaaagg	ccccctgctt	9060
gagtcctctt	cagggttagc	tctcagaaac	acaatctgca	gaacagattt	ttgttccaac	9120
atccttgcat	gagaatttgc	ccttagcttc	ccccacccca	gccaggctga	ataaaaattat	9180
gctgaaacta	ctgtcttatt	tgaggaaagt	aattagtcac	aggtggggag	gggtggggag	9240
attgcagaag	aatgttcatg	aattattagga	ttttcagctc	taagggggga	ctttgtaaac	9300
agctttagaa	gaagaaccag	gccggctggg	tgtggtggct	catgcctgta	atctcagcat	9360

ttggggaggc	caaggcgggc	ggatcacttg	aggtcaggag	tttgagacca	gcctggccaa	9420
catggtgaaa	ccctgtctct	attaaaaata	caaaaattag	ccagccgtgg	tagcgagcgc	9480
ctatgatccc	agctaactcc	gaggtctgag	ccagagaatc	acatgaacct	gggaggtgga	9540
ggctgcagtg	agccgagatc	acgccactgc	actccagcct	gggggacaga	gcaagaatct	9600
gtttcaaaaa	aaaaaaaaaga	aaaataggaa	ggaaggaaag	aaaggaaagg	aaagaagaga	9660
gagagaaaaga	aagagagaga	gagagagaga	aagaaaagaaa	gaaagaaaaga	aagaaaagaaa	9720
gaaagaaaaga	aagaaaagaaa	gaaaaagaaa	ggaaagaaaag	aacgaacgaa	ccaggcctcc	9780
ctctccaacc	ttcacctccg	tccctattct	ggccacttga	ttcgggggac	acctggtagg	9840
ggatggggaa	aggtgggagc	tgccagccag	aggggacccc	ggcttgagca	gcctcttgct	9900
gctatctgca	ggtttctcct	ggagactgaa	gtcgacctca	ggaagcccct	agagaacctg	9960
ggaatgaccg	acatgttcag	acagtttcag	gctgacttca	cgagtctttc	aggtaagaag	10020
actttccttt	gcattttctc	accccagtgg	actgcggggg	cccctaagag	gaaaaaggaa	10080
cctctccttg	agagcggcag	ctgatctaata	cctgtatcca	catctgtttc	agaccaagag	10140
cctctccacg	tgcgcagggc	gctgcagaaa	gtgaagatcg	aggtgaacga	gagtggcacg	10200
gtggcctcct	catccacagg	tgagtctggc	tcaggtgagg	ctccacgggt	gtcgcctcca	10260
tgcaccttca	ggataactgg	tccccagacc	cggaaaggac	cccgcagccc	tctcggcaca	10320
gagcagctct	gtctgtgctc	agccatcacc	cactccccac	ctgtttctca	gcctggaaaa	10380
cgggcttggg	accatggaac	cctgtttcct	cgctgatgg	ctcctaagtt	ccctgactgt	10440
gaaaaggcct	cctaaagaaa	aaccacaagt	gttcccacag	tgggaagtaa	acttaagaaa	10500
catgcttata	aggctgggca	tgggtggctcc	cacctgtaat	cccagcgctt	tgggggacca	10560
aggcaggtgg	atcacttgag	gtcaggaatt	cgagaccagc	ctgggcaaca	tggcaaaacc	10620
ctatctctac	taaaaataca	aaaattaggg	aggcgtgggtg	gcatgtgcct	gtagtcccag	10680
ctacttggga	ggctgaggca	ggagaatcac	ttgaatccag	gaggcagagg	ttgcagtga	10740
ccgagatcga	gctgctgcac	tccagcctgg	ccaatatagag	atgactctga	agaaaagaaa	10800
gaaagaaaaga	gagagagaga	gaaaagaaaag	aaagaaaagaa	agaaaagaaa	aaagaaaagaa	10860
agaaaagaaag	aaagaaaagaa	agagaaaagaa	agagaaaagaa	agaaaagaaa	gagcttatca	10920
ataagccctt	aaaggattta	gataaatgtg	tgtaaggga	gagctgatcc	attgctacca	10980
agctcctgga	ggaaaccagg	tctcagagga	tgtccctaaa	cttttaaggt	tcatattcag	11040
gaaaacaaac	aacttccagc	tgggcttagt	ggctcacacc	tgtaatccca	gcactttggg	11100
aggccgaggg	aggaggatcg	cttgagccca	ggaattttgag	accagcctgg	gcaatataat	11160
gagactgtgc	tctacaaaaa	ttagaaaaaa	attagccagg	catggtggca	tgcacctgta	11220
gccccagtta	cttgggagac	tgaggtggga	ggatcacttg	agcccatgag	ttcaaggctg	11280
cagtgagcca	tgaagggtgc	actgcactcc	cgctggggcg	acagagggag	accctgtctc	11340
taagaaaaac	ggcgggggtg	gggggtgggtg	cagtgccagc	atccctctgt	tctaagacat	11400
tgtcccttct	cttgagcgtg	tcatagtctc	agcccgcatg	gcccccgagg	agatcatcat	11460
ggacagaccc	ttcctctttg	tgggtccggca	caacccca	ggtgagcctg	gaacccatca	11520
cgttccacat	cctcccaccc	attctttctc	tcaggaacta	gtcccagacag	atgcagacat	11580
ccctctatcc	ctgagagggc	tctgggcagg	gaaccataac	cctaccctgc	ttcctgtccc	11640
aagaggaggc	taccttctat	caccacacaga	cagtgccggg	tccccgctct	gtgactcagg	11700
cagctgcgac	tccagacagc	tcaactcatct	gctatagatc	cagtccttcc	accacatcc	11760
agcctgatga	gctgtcccac	tccttctgct	ttcaacccc	catggttctt	ccacctcag	11820
gaacagtcct	tttcatgggc	caagtgtagg	aacctgacc	ctggggaaag	acgccttcac	11880
ctgggacaaa	actggagatg	catcgggaaa	gaagaaactc	cgaagaaaag	aatttttagtg	11940
ttaatgactc	tttctgaagg	aagagaagac	atttgccctt	tggtaaaaga	tggtaaacca	12000
gatctgtctc	caagaccttg	gcctctcctt	ggaggacctt	taggtcaaac	tccctagtct	12060
ccacctgaga	ccctgggaga	gaagtttgaa	gcacaactcc	cttaagggtct	ccaaaccaga	12120
cgggtgacgcc	tgcgggacca	tctggggcac	ctgcttccac	ccgtctctct	gcccactcgg	12180
gtctgcagac	ctggttccca	ctgaggccct	ttgcaggatg	gaactacggg	gcttacagga	12240
gcttttgtgt	gcttggtaga	aactatttct	gttccagtgca	cattgccatc	actcttgtac	12300
tgctggccac	cgcggaggag	gctgggtgaca	ggccaaaggc	cagtggaaga	aacacctttt	12360
catctcagag	tccactgtgg	cactggccac	ccctccccag	tacaggggtg	ctgcaggtgg	12420
cagagtgaat	gtcccccatc	atgtggccca	actctcctgg	cctggccatc	tccctcccca	12480
gaaacagtg	gcatgggtta	ttttggagtg	taggtgactt	gtttactcat	tgaagcagat	12540
ttctgcttcc	ttttattttt	ataggaatag	aggaagaaat	gtcagatgcg	tgcccagctc	12600
ttcaccccc	aatctcttgg	tggggagggg	tgtacctaaa	tatttatcat	atccttgccc	12660
ttgagtgcct	gttagagaga	aagagaacta	ctaaggaaaa	taatattatt	taaactcgct	12720
cctagtgttt	ctttgtggtc	tgtgtcaccc	tatctcagga	agtccagcca	cttgactggc	12780
acacaccctt	ccggacatcc	agcgtgacgg	agccacactc	gccaccttgt	ggccgcctga	12840
gaccctcgcg	ccccccgcgc	ccccccgcgc	cctctttttc	cccttgatgg	aaattgacca	12900
tacaatttca	tcctccttca	ggggatcaaa	gctcggaggt	ggggggacag	agactcagat	12960
gaggacagag	tggtttccaa	tgtgttcaat	agatttagga	gcagaaatgc	aaggggctgc	13020

```

atgacctacc aggacagaac ttcccccaat tacaggggtga ctcacagccg cattggtgac 13080
tcacttcaat gtgtcatttc cggctgctgt gtgtgagcag tggacacgtg aggggggggg 13140
tgggtgagag agacaggcag ctccgattca actaccttag ataatatattc tgaaaaccta 13200
ccagccagag ggtaggggcac aaagatggat gtaatgcact ttgggaggcc aaggcgggag 13260
gattgcttga gcccaggagt tcaagaccag cctgggcaac ataccaagac ccccgctctct 13320
ttaaaaaatat ataatattta aatatactta aatatatatt tctaatatct ttaaatatat 13380
atatatatatt taaagaccaa tttatgggag aattgcacac agatgtgaaa tgaatgtaat 13440
ctaatagaag cctaatacag ccaccatgtt ctccactgaa aaatcctctt tctttggggg 13500
ttttctttct ttcttttttg attttgcact ggacgggtgac gtcagccatg tacaggatcc 13560
acaggggtgg tgtcaaatgc tattgaaatt gtgttgaatt gtatgctttt tcacttttga 13620
taaataaaca tgtaaaaatg tttaaaaaaa ataataaaat aaataaaatc gaagaatatg 13680
tcaggacagt cactgccttc accttctcca ttccacaccg gtggtacaag aaatcagaag 13740
cctaggccag gtgtgggtgt tcatgcctgt aatcccagca ctttgggaag ccgagggtgg 13800
tggatcacct aaggtcagga gtttgagacc agcctggaca acatggtgaa acccgtctc 13860
tactaaaaat acaaaaatta gccgggcgtg gtggctggcg cctgtaatcc cagctactcg 13920
ggaggctgag gcaggagaat cacttgaagc caggaggcag aggttgcagt gagctgagat 13980
tgcaccactg aactccaggc tgggtggcag agcgagactc cctctcaaaa aacaacaact 14040
acaaagacaa caacaaaccc agaatacaaaa tcctgttggg ccatagacct catgggtgga 14100
agagaccttc ctacatccag gttggcccaa catgggggag tccatgaaat ggtcacctca 14160
gctctgccac aagccccaag gataagtga ttctgccctt gggaatcatc ctcaaaaagg 14220
aaaaaaatgt tccctgccca taaactttcc acttatgcag atgggcctgc tcgtaagtca 14280
ctgtcactgt ggggttcccaa ctctgttcat gacacttctt tccagcacca aatgcttccc 14340
acccctctac tccactccc cattcttcaa acccagctca agttccagtt cctccacctc 14400
ggacttccca tggatccagc caatatcact ctccaggtccg gcgcagtggc ccacgcctgt 14460
aatctcagca ctttgggagg ccggggcagg aagattgctt gaggccagga gtttcagacc 14520
agcctggaca acatagttag actc 14544

```

<210> 2
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 2
 aggaagaaat gtcagatgcg 20

<210> 3
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 3
 aggaatagag gaagaaatgt 20

<210> 4
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 4
 tgtcagatgc gtgccagct 20

<210> 5
 <211> 30
 <212> DNA
 <213> Homo sapiens

<400> 5
 aggaagaaat gtcagatgcg tgcccagctc 30

<210> 6
<211> 20
<212> DNA
<213> Homo sapiens

<400> 6
ataggaatag aggaagaaat 20

<210> 7
<211> 30
<212> DNA
<213> Homo sapiens

<400> 7
ttttattttt ataggaatag aggaagaaat 30

<210> 8
<211> 40
<212> DNA
<213> Homo sapiens

<400> 8
ttctgcttcc ttttattttt ataggaatag aggaagaaat 40

<210> 9
<211> 60
<212> DNA
<213> Homo sapiens

<400> 9
ttctgcttcc ttttattttt ataggaatag aggaagaaat gtcagatgcg tgcccagctc 60

<210> 10
<211> 16
<212> DNA
<213> Homo sapiens

<400> 10
gaagaaatgt cagatg 16

<210> 11
<211> 12
<212> DNA
<213> Homo sapiens

<400> 11
aagaaatgtc ag 12

INTERNATIONAL SEARCH REPORT

International Application No.

PO A2004/000424

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/53098 A (BILITEWSKI URSULA ; BIOTECHNOLOG FORSCHUNG GMBH (DE); KUHLMEIER DIRK ()) 26 November 1998 (1998-11-26) claims 12,13 page 2, paragraph 2 abstract ----- -/--	20-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

9 July 2004

Date of mailing of the international search report

22/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bellmann, A

INTERNATIONAL SEARCH REPORT

 International Application No
 PC 2004/000424

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BENZA: "Association of a Gene Polymorphism for PAI-1 with Primary Pulmonary Hypertension" CARDIOPULMONARY CRITICAL CARE NEWSLETTER, 'Online! XP002287063 Retrieved from the Internet: URL: http://www.americanheart.org/downloadable/heart/1017868019324CPCCSprng02.pdf 'retrieved on 2002-03-01!	18,19
A	page 8, column 2 - page 9, column 3 page 9, column 3	1-5,7, 9-17, 20-28
Y	HERMANS PETER W M ET AL: "4G/5G promoter polymorphism in the plasminogen-activator-inhibitor-1 gene and outcome of meningococcal disease" LANCET (NORTH AMERICAN EDITION), vol. 354, no. 9178, 14 August 1999 (1999-08-14), pages 556-560, XP002287064 ISSN: 0099-5355	18,19
A	page 556, column 1	1-5,7, 9-17, 20-28
Y	CATCHPOOLE D R ET AL: "The potential tumour suppressor role for caspase-9 (CASP9) in the childhood malignancy, neuroblastoma" EUROPEAN JOURNAL OF CANCER, PERGAMON PRESS, OXFORD, GB, vol. 37, no. 17, 2001, pages 2217-2221, XP004309829 ISSN: 0959-8049 abstract	18,19
Y	NOGUCHI E ET AL: "IDENTIFICATION OF MISSENSE MUTATION IN THE IL12B GENE: LACK OF ASSOCIATION BETWEEN IL12B POLYMORPHISMS AND ASTHMA AND ALLERGIC RHINITIS IN THE JAPANESE POPULATION" GENES AND IMMUNITY, NATURE PUBLISHING GROUP, GB, vol. 2, 2001, pages 401-403, XP002908314 ISSN: 1466-4879 the whole document ----- -/--	18,19

INTERNATIONAL SEARCH REPORT

International Application No

PO 2004/000424

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>GAILLARD M-C ET AL: "Identification of DNA polymorphisms associated with the V type alpha1-antitrypsin gene" BIOCHIMICA ET BIOPHYSICA ACTA . GENE STRUCTURE AND EXPRESSION, ELSEVIER, AMSTERDAM, NL, vol. 1444, no. 2, 16 February 1999 (1999-02-16), pages 166-170, XP004275307 ISSN: 0167-4781 the whole document</p>	18,19
A	<p>DAWSON SALLY J ET AL: "The two allele sequences of a common polymorphism in the promoter of the plasminogen activator inhibitor-1 (PAI-1) gene respond differently to interleukin-1 in HepG2 cells" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 15, 1993, pages 10739-10745, XP002287069 ISSN: 0021-9258 the whole document</p>	
A	<p>US 2002/086838 A1 (CHO SEONG H ET AL) 4 July 2002 (2002-07-04) the whole document</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2004/000424

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 6, 8
because they relate to subject matter not required to be searched by this Authority, namely:
Rule 39.1(iv) PCT - Diagnostic method practised on the human or animal body
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PO A2004/000424

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9853098	A	26-11-1998	AU 737771 B2	30-08-2001
			AU 4302297 A	11-12-1998
			CA 2290510 A1	26-11-1998
			WO 9853098 A1	26-11-1998
			EP 0983379 A1	08-03-2000
			JP 2001525678 T	11-12-2001
			RU 2193069 C2	20-11-2002
<hr/>				
US 2002086838	A1	04-07-2002	AU 2944101 A	24-07-2001
			WO 0151085 A1	19-07-2001
<hr/>				

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.